

STUDY ON THE ECOLOGY OF *Chenopodium murale* L. IN NILE DELTA, EGYPT.

Samia A. Haroun ; Ghada A. El-Sherbeny and Riyam S. Jasim
Department of botany, Faculty of Science, Mansoura University.

ABSTRACT

This study offers a description of the vegetation of *Chenopodium murale* community and evaluating the plant and soil parameters relationships. Classification analysis (TWINSPAN) revealed three vegetation groups; group A dominated by *Chenopodium murale*, group B dominated by *Melilotus indicus* and group C was dominated by *Pseudognaphalium luteo-album*. The ordination analysis (CCA) evaluate the *Chenopodium murale* and soil parameters relationship. The results indicated that *Chenopodium murale* showed a close relationship with soil clay, porosity, moisture content and total phosphorous whereas negatively correlated with electric conductivity, total dissolved salts, sodium and sulphate.

INTRODUCTION

Weeds are considered as an annoyance, especially in crop fields. *Chenopodium murale* is widely distributed weed which affect the crop productivity (Parker, 1997a).

Chenopodium murale belongs to family Chenopodiaceae. It is native to temperate Asia, Europe and Africa (Grin, 2000). In Egypt, it called Abu efein, Al zorbaih and Sonetar (Täckholm, 1974). It is described as an erect, grayish, annual weed, covered with mealy hairs, stem have few branches. The leaves alternating, broadly ovate to diamond-shaped, dentate. Inflorescence small, axillary and terminal paniculate spikes. Concerning distribution it grows at roadsides, gardens, damp habitats and canal banks (Felger, 2000 and Parker, 1997b). It grows also in new reclaimed land which may be salt-affected. Plant is used as a diuretic, anthelmintic, mild laxative, activating for liver, calmative. Leaves are edible as in U.A.E leaves used as a green salad. It is used as food in Africa and India (Holm *et al.*, 1997). Qasem and Abu-Blan (1995) reported that *C. murale* has a pesticidal effect and affected a wide range of microorganisms.

MATERIALS AND METHODS

The field studies were carried out during March and April 2014. 15 sites in the north-eastern Nile Delta of Egypt were selected for this study. In each site, all plants were documented in four plots (5x5 m²) and then summarized in one plot (100 m²). The frequency of each species was estimated at each site (Muller-Dombois and Ellenberg, 1974). The identification and nomenclature of plant species was, according to Täckholm (1974) and Boulos (1995, 1999, 2000, 2002, 2005, 2009).

Soil samples were collected at 0-50 cm depth at each site. Porosity, water holding capacity (WHC) and soil texture were estimated according to Allen *et al.* (1986). Organic carbon (OC) was measured by Walkely and Black's rapid titration. Soil water extract of 1:5 was used for estimation of conductivity ($\mu\text{S}/\text{cm}$), total dissolved salts (mg/L) and pH using a multi-meter CONSORT Model C535, while chloride (Cl^-) was estimated by direct titration against silver nitrate solution (N/35.5) and 5% potassium chromate indicator according to Jackson (1962). Carbonates (CO_3^{2-}) and bicarbonates (HCO_3^-) were estimated by titration with H_2SO_4 (0.1N) and phenol phthalein and methyl orange indicators as reported by Allen *et al.* (1986). Sulphates (SO_4^{2-}) were estimated using 5% barium chloride solution, (Piper, 1947). The classification of species data were statistically analyzed using the Two- Way Indicator Species Analysis TWINSpan (Hill, 1979 a&b).

RESULTS

The results of Two – Way Indicator Species Analysis (TWINSpan) illustrated in Figure (1) based on frequency values given three vegetation groups that reproduce the structure of the normal plant communities in the study area. Each group includes a set of sites that most homogenous of their vegetation from the other group. The data are given in Table (1).

Group A was the largest group, where it comprised eight sites situated in the relatively saline soils. This group was dominated by *Chenopodium murale* (F=62.4), while the indicator species include *Conyza bonariensis* (F=12.5). The other important species was *Cynodon dactylon* (F=45.23). Group B included four sites, this group was dominated by *Melilotus indicus* (F=50). In this group, the indicator species was *Coronopus didymus* (F=16.85) and the other preferential species were *Cakile maritime* (F= 50) and *Cynodon dactylon* (F= 50).

Group C involves three sites, this group was dominated by *Pseudognaphalium luteo-album* (F=66.67). The indicator species in this group was *Cynodon dactylon* (F=50). The most important species were *Paspalum distichum* (F=58.33) and *Solanum nigrum* (F=58.33).

Concerning the soil analysis, Table 2 showed the average values of soil properties of the different vegetation groups from TWINSpan analysis. The results indicated that group A revealed the lowest mean values of electrical conductivity (EC), total dissolved salts (TDS), bicarbonates (HCO_3^-), chlorides (Cl^-), sulphates (SO_4^{2-}), and clay (926.88 $\mu\text{S}/\text{cm}$, 474.75 mg/l , 0.07, 0.11 mg/l , 0.12 mg/l and 3.80% respectively). Meanwhile it possessed the highest value of organic carbon (OC= 1.05%), total phosphorous (TP= 8.09ppm), total nitrogen (TN= 18.02ppm), porosity (38.52%) and water holding capacity (50.40%).

Group B showed the highest mean values of pH (7.54), EC (1959 $\mu\text{S}/\text{cm}$), TDS (1075.25 mg/L) and sand (91.12%). Group C categorized by the second level of the mean value of EC (1846 $\mu\text{S}/\text{cm}$), TDS (958.67 mg/L) and MC (4.08%). This group attained the highest mean values of Cl^- (0.20 %) and the lowest mean values of TN (11.97ppm), TP (5.01ppm), porosity (30.70%) and sand (89.69%).

Figure (1): TWINSpan analysis of the 15 sites based on frequency Indicator species are abbreviate to the first three letters of the genus and the first three letters of species name respectively see Table 1.

Table (1): Frequency of *C. murale* and the associated species of different vegetation groups from TWINSPAN in the study area.

| | Taxon | A | B | C |
|----|--|-------|-------|-------|
| 1 | <i>Alternanthera sessilis</i> (L.)DC. | 10.25 | 9.25 | 8.00 |
| 2 | <i>Amaranthus lividus</i> L. | 0 | 6.25 | 0 |
| 3 | <i>Amaranthus hybridus</i> L. | 18.75 | 6.25 | 16.67 |
| 4 | <i>Symphytotrichum squamatum</i> (Spreng.) Nesom | 18.75 | 0 | 0 |
| 5 | <i>Atriplex portulacoides</i> L. | 25 | 25 | 0 |
| 6 | <i>Anagallis arvensis</i> var. <i>arvensis</i> L. | 0 | 0 | 0 |
| 7 | <i>Apium graveolens</i> (L.) Lag. | 0 | 0 | 16.67 |
| 8 | <i>Alhagi graecorum</i> Boiss. | 0 | 12.5 | 0 |
| 9 | <i>Bassia indica</i> (Wight) A.J.Scott. | 0 | 0 | 0 |
| 10 | <i>Beta vulgaris</i> L. | 18.75 | 6.25 | 0 |
| 11 | <i>Bidens pilosa</i> (L.) Lam | 0 | 0 | 0 |
| 12 | <i>Chenopodium ficifolium</i> Sm. | 12.5 | 6.25 | 0 |
| 13 | <i>Chenopodium album</i> L. | 0 | 0 | 8.33 |
| 14 | <i>Chenopodium glaucum</i> L. | 6.25 | 0 | 0 |
| 15 | <i>Chenopodium murale</i> L. | 62.4 | 37.5 | 50.00 |
| 16 | <i>Coronopus didymus</i> (L.) Sm. | 12.5 | 16.85 | 0 |
| 17 | <i>Cynodon dactylon</i> (L.)Pers. | 45.23 | 50 | 50.00 |
| 18 | <i>Cyperus rotundus</i> L. | 12.5 | 9.25 | 0.00 |
| 19 | <i>Cakile maritima</i> Scop. subsp. <i>aegyptiaca</i> (Willd.) Nyman | 0 | 50 | 16.67 |
| 20 | <i>Conyza bonariensis</i> (L.) Cronquist. | 12.5 | 0 | 0 |
| 21 | <i>Capsella bursae</i> Pastoris (L.) Medik. | 12.5 | 12.5 | 0 |
| 22 | <i>Convolvulus arvensis</i> L. | 25 | 37.5 | 0.00 |
| 23 | <i>Coronopus squamatus</i> (Forssk.) Asch. | 0 | 0 | 0 |
| 24 | <i>Cressa cretica</i> L. | 0 | 0 | 0 |
| 25 | <i>Euphorbia peplus</i> L. | 0 | 0 | 0 |
| 26 | <i>Erucaria hispanica</i> (L.) Druce | 0 | 0 | 0 |
| 27 | <i>Heliotropium curassavicum</i> L. | 0 | 6.25 | 0 |
| 28 | <i>Imperata cylindrica</i> (L.) Raeusch. | 25 | 0 | 8.33 |
| 29 | <i>Lepidium sativum</i> L. | 0 | 18.75 | 50.00 |
| 30 | <i>Limbarda crithmoides</i> L. Dumort. | 0 | 0 | 25.00 |
| 31 | <i>Lolium perenne</i> L. | 0 | 0 | 16.67 |
| 32 | <i>Malva parviflora</i> L. | 0 | 6.25 | 41.67 |
| 33 | <i>Melilotus indicus</i> (L.) All. | 0 | 50 | 16.67 |
| 34 | <i>Mentha longifolia</i> (L.) Huds. | 0 | 18.75 | 25.00 |
| 35 | <i>Mesembryanthemum crystallinum</i> L. | 0 | 0 | 0 |
| 36 | <i>Paspalidium gemmatum</i> (Forssk.) Stapf | 0 | 18.75 | 0 |
| 37 | <i>Paspalum distichum</i> L. | 0 | 12.5 | 58.33 |
| 38 | <i>Parapholis incurve</i> (L.) C.E. Hubb | 0 | 0 | 0 |
| 39 | <i>Persicaria salicifolia</i> (Willd.) Assenov | 0 | 12.5 | 0 |
| 40 | <i>Phragmites australis</i> (Cav.) Trin. exsteud. | 0 | 6.25 | 0 |
| 41 | <i>Plantago major</i> L. | 0 | 25 | 50.00 |
| 42 | <i>Polypogon monspeliensis</i> (L.)Desf. | 6.25 | 31.25 | 16.67 |
| 43 | <i>Portulaca oleracea</i> L. | 0 | 18.75 | 0 |
| 44 | <i>Pseudognaphalium luteo-album</i> (L.) Hilliard & B.L.Burt | 0 | 0 | 66.67 |
| 45 | <i>Ranunculus sceleratus</i> L. | 0 | 0 | 16.67 |
| 46 | <i>Rorippa palustris</i> (L.) Besser | 0 | 12.5 | 0 |
| 47 | <i>Rumex dentatus</i> L. | 0 | 12.5 | 0 |
| 48 | <i>Schismus barbatus</i> (L.) Thell. | 0 | 0 | 0 |
| 49 | <i>Senecio glaucus</i> L. | 0 | 0 | 16.67 |
| 50 | <i>Sisymbrium irio</i> L. | 12.5 | 25 | 8.33 |
| 51 | <i>Solanum nigrum</i> L. | 12.5 | 12.5 | 58.33 |
| 52 | <i>Sonchus oleraceus</i> L. | 0 | 0 | 0 |
| 53 | <i>Spergularia marina</i> (L.)Griseb. | 0 | 31.25 | 0 |
| 54 | <i>Suaeda maritima</i> (L.)Dumort | 25 | 37.5 | 50.00 |
| 55 | <i>Trifolium resupinatum</i> L. | 0 | 0 | 0 |
| 56 | <i>Urtica urens</i> L. | 0 | 0 | 0 |
| 57 | <i>Veronica anagallis-aquatica</i> L. | 0 | 0 | 0 |
| 58 | <i>Vicia sativa</i> L. | 12.5 | 0 | 0 |

Table (2) :Soil properties of the different vegetation groups from TWINSPAN in the study area.

| parameters | A | B | C |
|---|--------|---------|---------|
| pH | 7.09 | 7.54 | 7.06 |
| Electrical Conductivity $\mu\text{S}/\text{cm}$ | 926.88 | 1959.00 | 1846.00 |
| Total dissolved salts(mg/L) | 474.75 | 1075.25 | 958.67 |
| Bicarbonates % | 0.07 | 0.11 | 0.11 |
| Chlorides % | 0.11 | 0.15 | 0.20 |
| Sulphates% | 0.12 | 0.23 | 0.21 |
| Organic carbon% | 1.05 | 0.94 | 0.67 |
| Total nitrogen ppm | 18.02 | 14.68 | 11.97 |
| Total phosphorus ppm | 8.09 | 5.71 | 5.01 |
| Porosity% | 38.52 | 35.38 | 30.70 |
| Water holding capacity% | 50.40 | 47.81 | 48.60 |
| Clay | 3.80 | 3.95 | 5.21 |
| Silt% | 5.87 | 4.70 | 4.91 |
| Sand% | 90.27 | 91.12 | 89.69 |
| Moisture content% | 5.90 | 3.19 | 4.08 |

Species – soil relationships: Canonical Correspondence (CCA) analysis gives the ordination diagram concerning the relationship between the plant species and soil parameters in the study area (Figure2). The points represent species and the arrows represent environmental variables of soil.

The correlation between vegetation and soil characteristics was shown in the ordination diagram produced by CCA of the biplot of species and environmental variables (Figure 2). From the illustrated data, it is clear that *Chenopodium murale* showed close relationship with soil clay, porosity, moisture content and total phosphorous on the upper right side of CCA diagram. On the other hand, it showed a reverse correlation with electric conductivity, total dissolved salts, sodium and sulphate.




Figure (2): CCA analysis showing the soil and species relationship. The species abbreviation listed in Table 1.

DISCUSSION

The weed vegetation considered as a progression of plant species in a habitat where ecological conflict is mainly due to the crop management system. The productivity analysis of this vegetation is mainly focused on identifying the agronomic factors responsible for its variations (Ferrari *et al.*, 1984). Several studies have described the effects agronomic factors on species ecology and biology (Mohler & Liebman, 1987; Ghersa and Holt, 1995).

Radosevich & Holt (1984) reported that weeds represent an important constituent of the environment. Weeds steadiness is significant in vision of the efforts to remove them, and permits greater attention. Weed species affect the crop yield where reduce the farm revenue and represents an economically problem (Fayed *et al.*, 1997).

Application of TWINSPLAN analysis revealed that the vegetation of *C. murale* community produced three groups, group A was the largest group (8 sites) and dominated by *Chenopodium murale*, while the indicator species include *Conyza bonariensis*. The other important species was *Cynodon dactylon*. Group B included four sites, this group was dominated by *Melilotus indicus* and indicated by *Coronopus didymus*; the other preferential species were *Cakile maritime* and *Cynodon dactylon*. Group C involves three sites, this group is dominated by *Pseudognaphalium luteo-album* and indicated by

Cynodon dactylon The most important species are *Paspalum distichum* and *Solanum nigrum*. The dominants and associated species reflect their habitats.

Chenopodium murale showed a close relationship with soil clay, porosity, moisture content and total phosphorous and a reverse correlation with electric conductivity, total dissolved salts, sodium and sulphate. Moreover, *C. murale* recorded as nutrients collector such as nitrogen, phosphorous, potassium and magnesium (Maliwal and Gupta, 1988; Qasem, 1992).

REFERENCES

- Allen, S.E., Grimshaw, H.M., Parkinson, J.A., Quarmby, C., Roberts, J. D. (1986) Chemical Analysis. In: Chapman, S. B. (ed.), Methods in Plant Ecology, 411–466. Blackwell Scientific Publications, Oxford.
- Boulos, L. (1999-2005) Flora of Egypt, (Vols. I-IV) Al Hadara publishing Cairo, Egypt.
- Fayed, M. T. B., El-Geddawy, I. H. and el-zeny, M. M. (1997) Influence of weed interference on growth, yield and quality of sugar beet. *J. Agric. Res.*, 77 (3) 1239-1249.
- Felger, R.S. (2000). Flora of the Gran Desierto and Rio Colorado of northwestern Mexico. The University of Arizona Press. Tucson, Arizona. 673 pp.
- Ferrari, C.; Speranza, M. and Catizone, P. (1984). Weed and Crop Management of Wheat in Northern Italy. International Symposium on Weed Biology and Systematics, 7: 411-420.
- Ghersa, C. M., HOLT, J. S. (1995) Using phenology prediction in weed management: a review. *Weed Res.* 35(6), 461-470.
- Grin (2000). Grin Taxonomy. United States Department of Agriculture, Agricultural Research Service, The Germplasm Resources Information Network (GRIN). Website: <http://www.ars-grin.gov/npgs/tax/index.html> then click on 'simple queries of species data' and search for plant species.
- Hill, M. O. (1979). TWINSpan- a FORTRAN Program for ranging multivariate data in an ordered two way table by classification of individual and attributes. Section of Ecology and Systematic. Cornell Univ., Ithaca, New York.
- Hill, M.O.; Bunce, R.G.H.; and Shaw, M.W. (1975). Indicator species analysis, divisive polytheistic method of classification and its application to a survey of native pinewoods in Scotland. *J. Ecol.* Vol. 63, pp. 597 – 613.
- Holm, L.G.; Doll, J.; Holm, E.; Pancho, J.V. and Herberger, J.P. (1997). World Weeds: Natural Histories and Distribution. New York, USA: John Wiley & Sons Inc.
- Jackson, M. L. (1962) Soil Chemical Analysis. Constable and Co .Ltd. London, 496 p.

- Maliwal, P.L. and Gupta, O.P. (1988). Effect of herbicides and phosphorus on nutrient uptake in fenugreek and associated weeds. *Indian Journal of Weed Science*, 20(3):48-54
- Mohler, C.L. and Liebman, M. (1987). Weed Productivity and Composition in Sole Crops and Intercrops of Barley and Field Pea. *Jour. of Applied Ecology*, 24: 685-699.
- Muller-Dombois, D. and Ellenberg, H. (1974). Aims and methods of vegetation ecology. John Wiley & Sons, New York.
- Parker, R. (1997a). Control of problem weeds. In: William R.D., D. Ball, T.L. Miller, R. Parker, J.P. Yenish, R.H. Callihan, C. Eberlein, G.A. Lee, and D.W. Morishita (compilers). *Pacific Northwest 1997 Weed Control Handbook*. Oregon State University, Corvallis, Oregon. p. 304 -342.
- Parker, R. (1997b). Agrichemicals and their properties. In: William R.D., D. Ball, T.L. Miller, R. Parker, J.P. Yenish, R.H. Callihan, C. Eberlein, G.A. Lee, and D.W. Morishita (compilers). *Pacific Northwest 1997 Weed Control Handbook*. Oregon State University, Corvallis, Oregon. p.12 - 27.
- Piper, C. S. (1947). *Soil and Plant Analysis*. Interscience Publishers, Inc. New York.
- Qasem, J.R. (1992). Nutrient accumulation by weeds and their associated vegetable crops. *Journal of Horticultural Science*, 67(2):189-195.
- Qasem, J.R. and Abu-Blan, H.A. (1995). Antifungal activity of aqueous extracts from some common weed species. *Annals of Applied Biology*, 127(1):215-219; 21.
- Radosevich, S.R. and Holt, J.S. (1984). *Weed Ecology*. John Wiley and Sons, New York.
- Täckholm, V. (1974) *Student flora of Egypt*. 2nd ed. Published by Cairo Univ. Cooper. Prin. Comp, 887 pp.

دراسة بيئية على نبات السننار بدلنا النيل بمصر
سامية على هارون ، غادة عبد الله الشربيني و ريام صبيح جاسم الشمري
قسم النبات - كلية العلوم - جامعة المنصورة

هذه الدراسة تقدم وصف للكساء النباتي لعشيرة نبات السننار (الزربيج) و تقييم العلاقة بين النبات وعوامل التربة أوضح تحليل التصنيف ثنائي الاتجاه (TWINSpan) أن هناك 3 مجموعات : مجموعة A وكان يسوها نبات السننار مجموعة B وكان يسودها نبات الحندقوق ومجموعة C وكان يسودها صابونه العفريت . بتحليل التطابق الكنسي (CCA) اتضح أن نبات السننار كان له علاقة طردية مع التربة الطينية والمسامية والرطوبة والفسفور وعلاقة عكسية من التوصيل الكهربى والأملاح الكلية الذائبة والصوديوم والفسفات .




Figure (1): TWINSpan analysis of the 15 sites based on frequency Indicator species are abbreviate to the first three letters of the genus and the first three letters of species name respectively see Table 1.