

**MCQs: All questions to be answered:**

- 1. Enophthalmos in one eye without previous injury is suspicious for:**
  - a. cavernous hemangioma
  - b. orbital cellulitis
  - c. metastatic breast carcinoma in a woman
  - d. early orbital inflammatory disease
- 2. Prominent corneal nerves may be seen in which condition?**
  - a. pulmonary adenocarcinoma
  - b. lattice corneal dystrophy type I
  - c. Posterior polymorphous dystrophy
  - d. macular corneal dystrophy
- 3. Which of the following drugs is associated with corneal verticillata?**
  - a. Metoprolol
  - b. Erythromycin
  - c. Chloroquine
  - d. Tetracycline
- 4. Traumatic hyphema should typically be treated with which medicines?**
  - a. aminocaproic acid to slow clot dissolution and prevent rebleeding
  - b. topical corticosteroids and cycloplegic agents
  - c. tissue plasminogen activator to hasten dissolution of the clot
  - d. intravenous corticosteroids to reduce the inflammation
- 5. What is a typical finding in the early stages of *Acanthamoeba* keratitis?**
  - a. hypopyon
  - b. ring infiltrate
  - c. radial perineuritis
  - d. corneal neovascularization
- 6. Which gland does *not* contribute to the aqueous layer of the tear film?**
  - a. Krause
  - b. Main lacrimal
  - c. Zeis
  - d. Wolfring
- 7. Which one of the following statements regarding retinitis pigmentosa is true?**
  - a. The X-linked form is least common but most disabling.
  - b. Signs and symptoms typically precede ERG abnormalities.
  - c. Retinal pigmentary changes in the midperiphery are always present.
  - d. The initial visual field defect is a ring scotoma.
- 8. Which one of the following statements regarding Coats disease is true?**
  - a. It has an autosomal dominant pattern of inheritance with variable penetrance.
  - b. It is usually bilateral.
  - c. Males are affected more frequently than females.
  - d. It is usually diagnosed before 2 years of age.