

IMPROVING THE PROPERTIES OF RAS CHEESE MADE FROM GOATS MILK

El-Desoci, W. I.

Fac. of Agric., El-Azhar University, Assiut, Egypt

ABSTRACT

The effect of heat treatment of goat's milk and cold storage of milk pasteurization with or without refrigerated storage on the properties and keeping quality of Ras cheese was studied. Chemical composition, yield, ripening indices and microbiological properties were examined. Pattern of fatty acids and organoleptic properties of the resultant Ras cheese showed that the moisture, fat, yield and pH value of the cheese produced from goats milk were higher, while lower acidity was detected. Also values of ripening indices were higher in goat's milk cheese. On the other hand, a noticeable increase of short chain fatty acids in goats milk Ras cheese than cow's milk particularly T_2 however Ras cheese made from goats milk T_2 obtained the highest scoring points as in cow's milk cheese.

INTRODUCTION

Ras cheese is the main national variety hard cheese in Egypt and made from cow's milk or mixture of cows and buffalos milk. The properties of cheese as well as its keeping quality is governed by the properties and quality of milk used for its manufacture. Recently, goat's milk has been given great attention because, the goat's milk has a higher nutritional value than cow's milk, and it is recommended by some pediatrician for feeding infant. In a previous study (El- Abd et al., 1992), concluded that use of a mixture of goat's and cow's milk (1: 1) in making Ras cheese gave promising results. Also El- Batawy et al., (1992) used three salting methods in Ras cheese made from a mixture of goat's and cow's milk (1: 1). The chemical composition of Egyptian Goats milk was reported by El- Alamy et al., (1990). Hiekal et al., studied that the change in gross chemical composition, free fatty acids and organoleptic properties of Ras cheese made from cow, goat and mixed milk (1:1 cow to goat). On the other hand, with the changing practices in the dairy industry, milk is sometimes stored for extended periods before processing. The most appropriate method is refrigerated storage, although if extended, might lead to serious problems Yazid (1990). However, the nutritional value and composition of goat's milk is similar to cow's milk. Rakshy and Hassan (1971) and Isatchenko (1981) reported that the size of fat globule milk is nearly to be half the size of that in cow's milk. On the other hand, the content of whey proteins in goat's milk is higher than that in cow's milk. This led to the investigation of refrigerated storage as a means to limit bacterial growth (Diaman, 1993).

Besides, the goats milk contain more digestible proteins and fast incidence of allergy to goat's milk. Also the size of casein micelles in goat's milk is smaller than that in cow's milk (Dariani and Speck, 1980 and Battadjeva, et al., 1985). However, the duration for rennet coagulation of goats milk is longer and the gels is softer compared with cow's milk. There

were some trials to use goat's milk in making semi hard and hard cheese to overcome the problems which faced the cheese making from goats milk by heat treatment (Banks et al., 1987) and (Remeuf, 1992). However, little work has been carried out on Ras cheese which is considered the most popular hard type cheese consumed by the Egypt consumers. Also there was no information about the effect of heat treatment and cold storage on technological and chemical properties on the hard or semi hard cheeses produced.

The aim of this investigation, therefore, was to study the effect of heat treatment and cold storage of goat's milk on the properties of Ras cheese.

MATERIALS AND METHODS

Fresh cow's and goat's milks were obtained from the herd of the Animal Production Research Institute, Arab El-Awamer, Assuet Branch. Commercial fine grade salt was obtained from the local market. Lactic acid starters (*Lactococcus lactis* subsp. *lactis* and *Lactobacillus delbruekii* subsp. *bulgaricus*) and calf rennet powder were obtained from Chr. Hansen's Laboratories, Copenhagen Denmark. All media used for microbiological analysis were obtained from Oxoid Company (Hampshire, England) and Biolife S.R.L. (Milan, Italy).

Ras cheese was made from fresh cow's milk (3.5% fat), as described by Hofi et. al, (1970). Goats milk (3.5% fat) was divided into two parts, the first part (T_1) was heated 75°C / 15 sec. and used in the manufacture of Ras cheese as described by Hofi et. al., (1970). The second part was heated 80°C / 10 sec. and cold then divided to three equal portions. Ras cheese was directly manufactured from the first portion (T_2), the second and third portions were held at 5–7°C for 24 and 48 hr. (T_3 and T_4) respectively, then made into Ras cheese as described above (Okasha, 2001). The resultant cheeses were salted and waxed then ripened at 14°±1°C and 80–85% relative humidity (RH) for three months during which it was analyzed periodically.

Milk and cheese samples of all treatment were chemically analysed for moisture, titratable acidity (T.A), fat, pH value, salt, total nitrogen (T.N), soluble nitrogen (S.N) and non protein nitrogen (N.P.N) contents according to Ling (1963). The total volatile fatty acids (T.V.F.A) in cheese samples were determined by the method of Kosikowski (1982). Milk fat was extracted from cheese according to Rosegotleb method as described by the AOAC (1975), fatty acids pattern was determined by Gas-liquid chromatography (GLC) as explained by Deeth et. Al., (1983).

At the same time, cheese samples were examined for the presence of certain microbial groups including total bacterial count (T.C) were determined as suggested by the American Public Health Association (A.P.H.A,1992), proteolytic bacterial counts as recommended by Harriagan and McCance (1976) and lipolytic bacterial count was detected as suggested by American Public Health Association (A.P.H.A, 1992).

All cheese samples were organoleptically evaluated when fresh and during ripening every month by five staff members of the dairy department using the following scheme flavour (50), body& texture (40) and appearance (10) points as suggested by Abdel-Fattah (1966).

The data of all experiments were presented as the means of triplicate analysis. Statistical analysis was carried out using analysis of variance (ANOVA) and the differences between means were tested using Duncants test as well as average computer proram (SPSS,1999) at P<0.05.

RESULTS AND DISCUSSION

Data illustrated in Table (1) show that the chemical composition of cows and goats milk. Slightly higher content of T.S, fat and T.A were noticed in goats milk than the cow's milk, which agreed with those reported by Meislahn (1985), Ashmawy (1977), El-Alamy et al., (1990) and Girgis, et al., (1999). Variations in cheese yield of varios treatment are given in the same Table. As can be seen from the data, control cheese had the lower yield when fresh (before and after salting) and end ripening, compared with the all other treatments. These results are in agreement with those obtained by El-Batawy et al., (1992), El-Demerdash (1996) and okasha (2001).

Table (1): Chemical composition of milk used in cheese making and yield of the resultant Ras cheese during ripening period.

Parameters measured	C	T ₁	T ₂	T ₃	T ₄	F-test
T.S	11.68a±0.39	12.29a±0.31	12.30a±0.40	12.29a±0.45	12.28a±0.16	0.498ns
Fat	3.19a±0.03	3.32a±0.04	3.32a±0.02	3.32a±0.03	3.32a±0.02	2.259ns
T.A	0.16b±0.006	0.17ab±0.002	0.17ab±0.003	0.18a±0.004	0.175a±0.003	4.678 *
pH	6.75a±0.18	6.71a±0.15	6.71a±0.11	6.70a±0.09	6.71a±0.14	0.039ns
T.P	3.67a±0.07	3.52b±0.05	3.52b±0.06	3.51b±0.06	3.51b±0.05	3.228ns
Lactose	4.19a±0.02	4.25a±0.03	4.24a±0.021	4.24a±0.02	4.25a±0.11	0.856ns
Ash	0.71b±0.004	0.76a±0.003	0.76a±0.002	0.76a±0.003	0.76a±0.006	33.41**
Yield						
Before salting	11.46c±0.21	11.72bc±0.22	12.55a±0.13	12.32ab±0.16	11.89abc±0.15	4.308 *
After salting	10.92b±0.36	11.51a±0.38	12.00a±0.50	11.84a±0.18	11.32a±0.41	1.224ns
(90) days	8.51b±0.27	8.57b±0.29	9.32a±0.31	9.11a±0.35	8.61b±0.61	1.822ns

Control (C): Cow's milk Pasteurization standard.

Treatment (T₁): Goat's milk heated to (75°C / 15 sec.)

Treatment (T₂): Goat's milk heated to 8 °C / 10 sec. without cold storage.

Treatment (T₃): Goat's milk heated to 80°C / 10 sec. and cold storage at 5-7°C for 24 hr.

Treatment (T₄): Goat's milk heated to 80°C / 10 sec. and cold storage at 5-7°C for 48 hr.

a, b and c within the same row with different superscripts differed significantly(p<0.05).

(ns): insignificant.

(*): P < 0.05

(**): P < 0.01

Table (2) shows the change in the moisture content of Ras cheese made from either cows or goat's milk and all treatment. The changes in

moisture content of fresh Ras cheese as affected by heat treatment of cheese milk were higher than that in the control. It is clear from these data that the moisture content of all cheeses decreased gradually throughout the ripening period. This might be due to some of the water gradually evaporates during ripening or becomes bound with the protein as about water, as ripening progressed (Conner, 1988). On the other hand, the cheese made from milk heated and cold stored for different period had higher moisture content than that made from heated without cold storage. Similar results were obtained by El-Attawy (1980), El-Abd et al., (1992), Okasha (2001) and Hefnawy (2004).

From results present in Table (2), it could be concluded that the fat content in dry matter (D.M) of Ras cheese of all treatments increased gradually with the advancing of the ripening period. Also all Ras cheeses made from goats milk had higher fat content than those made from goats milk. The general trend of those results was in accordance to those reported by Fikry (1987), Shendy (1989), Harby (1991), El- Batawy et al., (1992) and Okasha (2001).

Results given in Table (2) reveal the total nitrogen (T.N %) and T.N in D.M content of the different treatments of Ras cheese. It is clear that there was no considerable difference in T.N content between the treated cheeses (made from goat's milk) and control (mad from cow's milk). These results are in agreement with those obtained by El-Demerdash (1996), Guinee et al. (2000) and Okasha (2001).

Concerning the salt content and salt water ratio (S.W.R), results the same Table generally show that the average of S.W.R. in all treatment increased till the end of ripening period. On the other hand, the salt water ratio of the control were slightly higher than in all cheeses when fresh cheese. The results are in agreement with Harby (1991), El- Batawy et al., (1992) Grgis et al., (1999), Abo-El-Nor (2004) and awad (2006).

The titratable acidity (T.A) was higher in the control than the all treatments when fresh and during ripening period. On the other hand, the acidity of all cheese samples gradually increased with the increase of storage period probably as a result of fermentation of residual lactose and degradation of protein and fat (Hofi et al., 1991). The pH values rang parallel to the titratable acidity when fresh and throughout storage period. However, pH values of Ras cheese were decreased gradually in all treatments during ripening until 45 days while were increased until the end period 90 days. This is in agreement with that found by El-Abd et al., (1992), El-Demerdash (1996), Mostafa (1999) and Osman (2003).

Table (2): Effect of heat treatments and cold storage of Goats milk on chemical composition of Ras cheese during ripening period.

Ripening period (days)	C	T ₁	T ₂	T ₃	T ₄	F-test
Moisture%						
0	44.20b±1.18	45.27ab±1.21	47.15a±1.53	47.19a±1.14	46.51a±2.73	3.87ns
15	40.15a±1.17	41.23a±2.00	41.20a±1.72	41.85a±1.02	41.62a±1.67	0.68ns
30	36.71a±1.08	37.20a±1.16	37.81a±2.23	37.90a±1.17	37.95a±1.16	0.63ns
45	35.92a±1.15	36.21a±1.12	36.71a±1.13	36.20a±1.13	36.05a±1.77	0.19ns
60	33.21a±1.50	34.55a±0.88	35.20a±0.68	35.12a±1.08	35.72a±1.61	1.58ns
90	32.84a±1.08	32.10a±1.13	33.60a±1.02	34.10a±0.68	33.95a±1.07	1.83ns
Salt water ratio S.W.R%						
0	3.61a±0.36	3.49a±0.29	3.43a±0.29	3.48a±0.26	3.50a±0.31	0.53ns
15	5.28a±0.32	4.89ab±0.36	4.61ab±0.36	4.54b±0.41	4.73ab±0.36	2.01ns
30	6.02a±0.31	5.88ab±0.21	5.28b±0.21	5.25b±0.33	5.61ab±0.21	2.78ns
45	6.29a±0.30	6.18a±0.33	5.96a±0.45	6.10a±0.16	6.10a±0.22	0.22ns
60	6.95a±0.46	6.65a±0.31	6.39a±0.26	6.37a±0.27	6.35a±0.25	1.73ns
90	7.21a±0.37	7.28a±0.19	6.57bc±0.20	6.51bc±0.21	6.49bc±0.17	2.50ns
Fat /D.M%						
0	45.21a±0.75	46.34a±1.43	46.24a±2.63	46.42a±1.10	46.31a±0.69	0.33ns
15	49.51a±1.66	48.52a±1.18	48.64a±3.95	48.21a±0.68	50.44a±1.12	1.23ns
30	52.64a±1.6	50.66ab±0.68	52.11b±1.11	52.45a±0.61	50.11ab±2.16	2.78ns
45	52.78a±1.18	51.63a±2.12	52.57a±0.89	52.84a±1.31	51.83a±0.71	0.82ns
60	53.10a±2.23	52.15a±2.16	52.71a±1.23	53.19a±1.17	52.52a±1.08	0.28ns
90	53.75a±1.14	52.77a±1.21	53.92a±1.15	53.97a±1.22	52.71a±0.61	0.66ns
T.N %						
0	3.72a±0.09	3.83a±0.11	3.77a±0.11	3.92a±0.10	3.81a±0.10	0.47ns
15	4.05a±0.18	4.32a±0.13	4.11a±0.17	4.22a±0.10	4.14a±0.13	1.44ns
30	4.39a±0.11	4.47a±0.18	4.49a±0.13	4.44a±0.18	4.41a±0.15	0.09ns
45	4.51a±0.17	4.64a±0.10	4.69a±0.19	4.60a±0.13	4.56a±0.11	0.45ns
60	4.70a±0.09	4.77a±0.17	4.88a±0.10	4.83a±0.10	4.67a±0.11	0.67ns
90	5.15a±0.19	5.20a±0.11	5.17a±0.45	5.23a±0.23	5.11a±0.32	0.32ns
T.A						
0	1.07a±0.066	1.00a±0.06	0.95a±0.06	0.93a±0.06	0.92a±0.06	0.97ns
15	1.14a±0.11	1.10a±0.16	1.06a±0.11	1.01a±0.09	1.00a±0.09	0.69ns
30	1.37a±0.09	1.24ab±0.11	1.15b±0.06	1.12b±0.06	1.10b±0.11	3.06ns
45	1.50a±0.11	1.45a±0.09	1.41a±0.08	1.38a±0.11	1.32a±0.15	1.23ns
60	1.66a±0.06	1.62ab±0.10	1.57ab±0.04	1.50ab±0.06	1.42b±0.06	2.73ns
90	1.92a±0.03	1.85ab±0.06	1.77ab±0.06	1.73b±0.11	1.70b±0.11	2.82ns
pH values						
0	5.67a±0.16	5.68a±0.13	5.71a±0.20	5.75a±0.13	5.82a±0.13	0.29ns
15	5.51a±0.26	5.53a±0.22	5.59a±0.22	5.63a±0.12	5.70a±0.12	0.29ns
30	4.72a±0.22	4.77a±0.33	4.80a±0.13	4.85a±0.13	4.92a±0.16	0.57ns
45	4.61a±0.13	4.68a±0.13	4.72a±0.11	4.79a±0.22	4.83a±0.22	0.68ns
60	5.32b±0.13	5.95a±0.13	5.50b±0.13	5.45b±0.13	5.38b±0.13	3.87*
90	5.59a±0.22	5.69a±0.13	5.65a±0.21	5.71a±0.15	5.65a±0.46	0.22ns

See legand table (1)

Table (3) shows the changes in the soluble nitrogen (S.N), non protein nitrogen (N.P.N), S.N/T.N,N.P.N/T.N and total volatile fatty acids (T.V.F.A). It is clear from these results that the ripening indices increased

gradually during ripening period. Also those data revealed that using goat's milk in Ras cheese making showed high ratio of S.N/T.N% and NP.N/T.N% which means, more acceleration of cheese ripening than cow's milk. Also the T.V.F.A content of Ras cheese made from goat's milk was higher when compared with the control. On the other hand, it could be observed that the rate of ripening indices increased during ripening period were relatively higher in T₂ than that of control and the other all treatments. The general trend of those results agreed with those reported by Hiekal et al., (1992), El- Abd et al., (1992) Ashmawy (1997), Shendy (1998), Mostafa (1999), Girgis et al., (1999) and Okasha (2001).

Table (3): Effect of heat treatments and cold storage of Goats milk on ripening indices of Ras cheese during ripening period.

Ripening period (days)	C	T ₁	T ₂	T ₃	T ₄	F-test
S.N%						
0	0.15a±0.012	0.18a±0.02	0.17a±0.01	0.16a±0.01	0.17a±0.01	0.82 ns
15	0.20b±0.02	0.24ab±0.04	0.25a±0.02	0.23ab±0.01	0.22ab±0.01	2.19 ns
30	0.30b±0.03	0.35a±0.01	0.33ab±0.02	0.32ab±0.02	0.31b±0.02	2.78 ns
45	0.39c±0.01	0.48a±0.02	0.44b±0.01	0.42bc±0.01	0.41bc±0.03	8.78**
60	0.44b±0.04	0.51a±0.01	0.50a±0.012	0.47ab±0.01	0.46ab±0.014	2.73 ns
90	0.63b±0.01	0.67a±0.01	0.61b±0.02	0.60b±0.02	0.59b±0.01	6.48**
N.P.N%						
0	0.055a±0.03	0.063a±0.02	0.074a±0.02	0.065a±0.02	0.069a±0.02	0.04 ns
15	0.11a±0.04	0.15a±0.01	0.14a±0.04	0.13a±0.02	0.15a±0.04	0.57 ns
30	0.20a±0.02	0.22a±0.03	0.23a±0.02	0.25a±0.04	0.20a±0.03	0.99 ns
45	0.22a±0.02	0.24a±0.02	0.27a±0.02	0.27a±0.03	0.23a±0.02	1.12 ns
60	0.25b±0.02	0.27ab±0.01	0.29a±0.02	0.30a±0.04	0.27ab±0.02	3.87*
90	0.30a±0.03	0.34a±0.02	0.35a±0.04	0.32a±0.02	0.31a±0.02	0.75 ns
S.N./T.N%						
0	4.03a±0.37	4.69a±0.44	4.50a±0.44	4.08a±0.76	4.46a±0.74	0.42 ns
15	4.93a±0.76	5.25a±0.43	6.08a±0.76	5.45a±0.44	5.31a±0.71	0.93ns
30	6.83a±0.71	7.82a±0.76	7.34a±0.81	7.20a±0.44	7.02a±0.44	0.72ns
45	8.64b±0.40	10.34a±0.44	9.38ab±0.76	9.13ab±0.70	8.99ab±0.79	2.14ns
60	9.36b±0.76	10.69a±0.39	10.24ab±0.44	9.73ab±0.8	9.85ab±0.44	2.81ns
90	12.23ab±0.52	12.88a±0.76	11.69ab±0.73	11.47b±0.71	11.54ab±0.4	2.24ns
N.P.N/T.N%						
0	1.47a±0.21	1.64a±0.23	1.96a±0.17	1.65a±0.36	1.81a±0.31	0.97 ns
15	2.71b±0.36	3.47a±0.19	3.40a±0.21	3.08ab±0.21	3.62a±0.37	3.15 ns
30	4.55b±0.33	4.92b±0.36	5.12ab±0.36	5.63a±0.32	4.53b±0.21	4.86*
45	4.87c±0.24	5.71ab±0.41	5.75ab±0.36	5.86a±0.33	5.04bc±0.26	4.01*
60	5.31b±0.31	5.66ab±0.32	5.94ab±0.33	6.21a±0.31	5.78ab±0.21	0.73 ns
90	5.82b±0.39	6.53ab±0.35	6.76a±0.34	6.11ab±0.24	6.06b±0.23	3.24ns
T.V.F.A 0.1 NaOH/100g cheese						
0	10b±1.18	13a±0.66	12ab±0.98	11ab±1.32	12ab±1.63	2.29 ns
15	16c±1.36	20ab±0.79	19b±0.79	22a±0.79	21ab±0.79	8.60**
30	24b±1.26	29a±1.36	30a±0.71	26b±0.77	29a±0.1.36	10.21**
45	29c±0.98	35b±2.23	39a±1.62	32bc±1.69	34b±1.76	12.99**
60	35d±1.11	42ab±1.69	44a±0.79	40bc±0.79	39c±0.79	17.47**
90	40c±1.69	49a±2.23	43bc±1.32	44b±0.72	42bc±1.29	9.09**

See legand table (1)

The concentration of saturated free fatty acids and short chaine acids increased in Ras cheese mad from goat's milk as the ripening period than cow's milk cheese Table (4). On the other hand, results showed fatty acids, which play an important role in flavor formation of cheese Omar (1984). Also cold storage of heated milk for 24 hours (T_3) caused more increase in the cold storage period (T_4) more increase in the short chaine fatty acids was observed. These results are in agreement with Hagag et al., (1987), Hiekal et al., (1992), El-Demerdash (1996), Mostafa (1999) and Okasha (2001).

Table (4): Effect of heat treatment and cold storage of goats milk on fatty acids in Ras cheese during ripening period.

Treatments	Short chain fatty acids	Middle chain fatty acids	Long chain fatty acids	Total saturated fatty acids	Total unsaturated fatty acids
30 days					
C	4.66d \pm 0.49	42.22ab \pm 0.65	53.12a \pm 0.76	59.24b \pm 1.15	40.76a \pm 1.72
T_1	5.35cd \pm 0.35	40.23b \pm 1.13	54.42a \pm 1.19	61.72b \pm 1.67	38.28b \pm 0.96
T_2	6.75c \pm 0.81	43.51a \pm 1.09	49.74b \pm 0.84	68.83a \pm 2.31	31.17d \pm 0.78
T_3	21.40b \pm 0.85	40.15b1.13	38.45c \pm 0.86	67.13a \pm 1.65	32.87c \pm 0.85
T_4	25.82a \pm 0.67	36.71c \pm 1.16	37.47c \pm 0.81	66.23a \pm 2.17	33.78c \pm 1.04
F-tes	426.4**	15.74**	273.3**	12.97**	60.00**
60 days					
C	7.77c \pm 0.84	46.84b \pm 1.12	45.39a \pm 0.95	63.51c \pm 2.25	36.49a \pm 0.95
T_1	8.51c \pm 0.79	45.92b \pm 1.65	45.57a \pm 1.33	67.32b \pm 1.81	32.68b \pm 0.95
T_2	9.22c \pm 0.85	49.93a \pm 1.13	40.85b \pm 0.63	72.15a \pm 1.65	27.85d \pm 1.51
T_3	27.12b \pm 0.65	47.01b \pm 1.16	25.87d \pm 0.86	69.77ab \pm 2.17	30.23c1.34
T_4	30.50a \pm 1.13	40.61c \pm 1.11	28.89c \pm 0.9	70.11ab \pm 2.10	29.89c \pm 0.65
F-tes	388.2**	27.38**	371.9**	10.74**	26.95**
90 days					
C	9.21e \pm 0.85	47.11b \pm 0.88	43.68a \pm 1.01	69.18b \pm 1.67	30.82a \pm 0.80
T_1	11.11d \pm 0.53	46.07b \pm 0.65	42.82a \pm 0.85	71.11ab \pm 1.68	28.89b \pm 0.87
T_2	14.48c \pm 0.53	51.26a \pm 0.69	34.26b \pm 0.77	73.92a \pm 1.65	26.08c \pm 0.61
T_3	31.22b \pm 0.51	46.73b \pm 1.13	22.05d \pm 0.63	71.81ab \pm 2.25	28.19b \pm 0.45
T_4	33.14a \pm 0.81	38.51c \pm 0.66	28.35c \pm 0.98	71.62ab \pm 1.67	28.38hi \pm 0.95
F-tes	616.0**	66.48**	303.6**	2.74ns	13.31**

See legand table (1)

Data illustrated in Table (5) show that the total, proteolytic and lipolytic bacterial counts of all treatments during ripening period of Ras cheese. It could be seen that the T.C, P.B.C and L.B.C of the control were lower than in all treatments when fresh and until 45 days. On the other hand TC, P.B.C and L.B.C of cheese of all treatments gradually decreased until the end of ripening period. This conclusion was in agreement which El-Shafi and Farag (1994) and Okasha (2001).

Table (5): Effect of heat treatments and cold storage of goat's milk on certain microbiological properties of the resultant Ras cheese.

Ripening period (days)	C	T ₁	T ₂	T ₃	T ₄	F-tes
Total viable count x 10 ⁵ (cfu/g)						
0	201d±4.46	229a±3.31	222b±5.33	218bc±2.34	211c±3.23	24.28**
15	147c±3.46	191a±2.01	180b±2.99	187a±3.44	192a±3.45	101.3**
30	116d±3.35	137c±3.46	145b±3.46	151b±4.21	162a±4.41	74.60**
45	99b±3.42	111a±4.11	92b±2.76	95b±3.76	96b±2.15	13.70**
60	60bc±3.73	81a±4.79	56cd±3.42	52d±3.31	64b±1.99	24.31**
90	37b±3.52	41ab±2.16	35b±3.41	44a±1.32	47a±3.23	5.84**
Proteolytic bacterial count x10 ⁴ (cfu/g)						
0	125c±2.13	149a±1.53	114e±2.59	121d±1.13	145b±1.13	409.4**
15	92c±1.06	98b±1.13	91c±2.54	96b±1.33	105a±3.00	30.03**
30	81d±2.71	87b±1.31	82d±2.31	85c±1.66	90a±1.45	36.67**
45	62cd±1.13	66b±2.11	61d±0.65	65bc±2.70	71a±1.31	16.33**
60	45b±1.15	46b±0.65	40c±1.07	46b±1.44	51a±2.25	42.10**
90	34c±0.65	37b±0.65	33c±1.13	39b±1.07	42a±2.14	21.70**
Lipolytic bacterial count x10 ³ (cfu/g)						
0	103c±2.18	110b±0.87	101d±5.14	97e±0.86	114a±1.47	140.7**
15	77d±0.96	84b±1.48	73e±2.12	81c±1.55	94a±0.86	151.6**
30	70c±1.47	74b±0.86	61e±1.87	67d±1.82	80a±0.54	155.5**
45	60b±0.86	61b±1.47	48d±0.93	55c±0.81	64a±1.86	110.1**
60	42b±1.86	40c±0.54	35d±1.18	38c±0.96	49a±3.11	85.12**
90	23c±0.58	27b±0.52	21d±0.86	25c±1.47	33a±2.18	65.12**

See legand table (1)

Cheeses of all treatments were organoleptically evaluated when 30 days and each 30 days up to 3 months of ripening period. Cheese made from goat's milk was generally characterized by a smooth texture goatly flavour and more piquant taste than cheese made from cow's milk. Data presented in Table (6) show that the flavour developed more rabidly and reached higher and almost similar scores in both the control and T₂ (Ras cheese made from heating the goats milk at 80°C for 10 sec. without cold storage). It was followed by T₁, T₃ and T₄ in a descending order.

Table (6): Effect of heat treatments and cold storage of goat's milk on the organoleptic properties of the resultant Ras cheese during ripening period.

Month	Treatments	Flavoure (50) points	Body& texture(40) points	Appearance (10) points	Total scores (100) points
1	C	35a±0.76	28a±0.49	7a±0.049	70a±1.07
	T ₁	34ab±0.71	24c±0.48	6b±0.050	64bc±1.06
	T ₂	34ab±0.81	26b±0.42	7a±0.050	67ab±1.10
	T ₃	32bc±0.65	23cd±0.39	7a±0.030	62cd±1.20
	T ₄	31c±0.56	22cd±0.75	6b±0.040	59c±1.70
	F-tes	4.72*	24.08**	124.6**	16.04**
2	C	41a±0.85	33a±0.52	8a±0.031	82a±2.10
	T ₁	38bc±0.63	30bc±0.29	7b±0.100	75b±2.20
	T ₂	40ab±0.31	31b±0.22	8a±0.060	79a±0.90
	T ₃	37c±0.33	30bc±0.26	7b±0.020	74b±1.60
	T ₄	36c±0.37	29c±0.21	7b±0.060	72b±1.50
	F-tes	7.52**	9.55**	124.6**	14.29**
3	C	42a±0.65	35a±0.41	8a±0.035	85a±2.50
	T ₁	40a±0.73	31b±0.26	8a±0.090	79b±1.30
	T ₂	47a±0.42	35a±0.33	8a±0.040	81a±1.20
	T ₃	40a±0.34	30bc±0.27	8a±0.070	78b±1.10
	T ₄	40a±0.76	29c±0.22	8a±0.020	77b±1.30
	F-tes	1.40 ns	33.22**	0.001 ns	11.66**

See legand table (1)

REFERENCES

- American Public Health Association (A.P.H.A) (1992). Standard Methods for the Examination of Dairy Products. A.P.H.A, Inc., 12th ed, New York.
- A.O.A.C. (1975). Official Methods of Analysis (12 th edo.) Association of Official Analytical Chemists Washington, DC.
- Abd-EI -Fattah, E.A. (1966). Studies on butter milk and its utilization in the manufacture of baby Edam cheese. MSc. Thesis, Ain Shams Univ. Egypt.
- Abou- EI- Noure, A.M., Abbas, F.M., Khalil, R.A.M. and El- Safty, M, S. (2004). Modified manufacture procedure for improving the quality of low fat Ras cheese. Egyptian J. Dairy Sci, 32: 141-152.
- Ashmawy, T.A.M. (1997). Studies on some factors affecting yield, chemical and technological properties of milk on Baladi goats and their crosses. M. Sc. Thesis. Food Sci., Department Faculty of Agric. Zagazig Univ. Egypt.
- Awad, S. (2006). Texture and flavours development in Ras cheese mad from raw and pasteurized milk. Food chemistry, Vol. 97, Issue 3, 394-400.

- Banks, J.M., Stewart, G.D. and Wast, I.G. (1987). Increasing the yield of cheddar cheese by the acidification of milk containing heat denatured whey proteins. *Milchwissenschaft*, 42 (4) 212-215.
- Battadjeva, M., Kalatzopoulos, G. AND Stamenova, V. (1985). free fatty acids and amine acid in two cheese mad from goats milk. *Lait* 65: 650-659.
- Conner, T. (1980). Advances in accelerated in ripening of cheese. *Cultured Dairy Prod. J.* 11: 21- 25.
- Dariani, D.N. and Speck, S.J. (1980). Soft pickled cheese made from cows and goats milk. II. Consumer evaluation. *J. International Goat and Sheep Research* I. 210-215.
- Deeth, H.C., Fits-Gerald, C.H. and Snow, A. F. (1983). Ages chromatographic method for the quantitative determination of free fatty acids in milk and products. *N. Z. J. Dairy Sci., Technol.* 18. 13-20.
- Diaman, T.M. (1993). The effect of refrigeration conditions on the level of bacterial contamination of milk molanco- Myanse Skatartvo 83: 56. C.F. *Dairy Sci., Abst.* 57: 3033.
- EL-Alamy, h. A., Mahfouz, M.b., hamzawy, I.f. and El-Loly, M.M. (1990). Studies on the chemical composition of Egyptian goats milk. I. Gross chemical composition and mineral contents. *Egyptian J. Dairy Sci.* 18 : 195.
- El- Attawy, Z.A. (1980). The use of milk powder in hard cheese making Ph. D. Thesis Ain Shams Univ. Egypt.
- El-Abd,M.M., El-Battawy, M.A., Younes, N.A. and El- Tawel, H.S. (1992). Utilization of goats milk in making Ras cheese. *Egyptian J.Dairy Sci.*, 20: 91.
- El-Batawy, M.A., El-Abd, M.M. Younes, N.A. and El-Rawel, H.S. (1992). Effect of salting method on the ripening of Ras cheese made from mixed goats and cow's milk. *Egyptian J. Dairy Sci.*, 20: 341-350.
- El-Demerdash, M.E. (1996). Studies on Gouda cheese. Ph. D. Thesis. Zagazig Univ Egypt. Elein, S., Girgis, I.H.I., Abd -El-Ghany, Laili,M. Yousef, and Laila, m. m. Mohamed (1991). Effect of milk pre treatment and storage conditions on the properties and quality of Ras cheese. *Egyptian J. Dairy Sci.*, 27: 153- 166.
- El-Shafie, M.M. and Farag, A.A. (1994). Improving the quality of smoked Ras cheese *Egyptian J. Appl. Sci.*, (9): 400-411.
- Fikry, S.A. (1987). Chemical and bacteriological studies on Edam cheese. MSc Thesis Fac. Of Agric. Univ. Egypt.
- Guinee, T., Auty, M.A.E. and Fenelon, M.A. (2000). The effect of fat content on the rheology, microstructure and heat induced functional characteristics of cheddar cheese. *International Dairy J.* 10: 277.
- Hagag, H.F., Hamzawy, L.F. and Shahin, Y. (1987). Fatty acids composition of globule core lipids from Egyptian cow, buffaloes and goats milk. *Egyptian J. Dairy Sci.*, 15: 25.
- Harby, S.I. (1991). Microbiological studies on hard cheese .Ph. D. Thesis. Cairo Univ.
- Harrigan, W.F. and McCance, E. (1976). *Laboratory Methods in Food and Dairy Microbiology*. Academic Press, New York. U.S.A.

- Hefnawy, A.A., El-Alfy, M.P., Nasr, W.I. and Sonia M. Abdou (2004). Potential of making Ras cheese from milk concentrated by U.F. and recombination techniques. Egyptian J. Dairy Sci., 32: 163-173.
- Hiekal, A.H., Kandeel, A.A. and Shendy, A.M. (1992). Relationship between type of milk and free fatty acids content of Ras cheese. Egyptian J. Dairy Sci., 20: 53-65.
- Hofi, A.A., Abd-El Hamid, L.B., Ahmed, N.S. and Abbas H.M. (1991). Acceleration of Ras cheese ripening by relevant slurry. Egyptian J.Dairy Sci., 19 (2): 277-346.
- Hofi., A.A. Yousef, E.H. and Goninein, M.A. (1970). Keeping changes in Cephalotyre cheese manufacture from raw and pasteurized milk with special reference to flavour. J. of Dairy Sci., 53: 1205.
- Isatchenko, Nauka zizni 7-122-123 (1981). Jenness, R. (1980). Composition and characteristics of goats milk as distinct from those of cow's milk, IDF. Doe, No. 202 pp. 54-67.
- Kosikowski, F. (1982). Cheese and fermented milk Food. Second Ed. 3rd printing with revisions, F.V. Kosikowski and Associates P.O.Box. 139, Brooktontdsle New York, 14817.
- Ling, E.R. (1963). A text Book of Dairy chemistry. Vol. 2 Partical 3rd Ed.Champan and Hall London.
- Meslahen, H.D. (1985). Thermization for improving the quality of supplier milk stored for 3 days. Thesis Tieraztliche, Hanover, German Federal Republic P. 98. C.F. Dairy Sci., Abst. 49: 477.
- Mostafa, M.B.M. (1999). Quality and ripening of Gouda cheese from goats milk as affected by certain feeding rations. Egyptian J. Dairy Sci., 27: 179- 190.
- Okasha, I.I. (2001). Effect of heat treatment and cold storage of goat's milk on some properties of Ras cheese during ripening. Egyptian J. Dairy Sci. and Techno. 357- 368.
- Omar, M.M. (1984). Microstructure free amino acids and free fatty acids in Ras cheese. Food chemistry 15: 19.
- Osman, M.M. (2003). Acceleration of the ripening and flavour development of Ras cheese using *Brevibacterium linens*. Egyptian J. Dairy Sci., 31:159-172.
- Rakshy, S.E. and Hassan, N. (1971). The suitability of goat's milk for the manufacture of a pickled cheese variety. Mex. J. of Agric. Res. 19: 269- 272.
- Remeuf, F. (1992). International conference goats pre-conf, proc. Invited papers. Vol. 2, part 3, 374-381 Marthur, India central Inst Res. On goats.
- Shendy, A.Y. (1989). Studies on goats milk utilization in soft and hard cheeses making. Ph.D. Thesis. Al-Azhar Univ.
- SPSS (1991). SPSS for Windows. Release 10.0 Standard Version. Copyright SPSS Inc., 198- 1999.

Yazid, B.H.A.MM. (1990). Effect of low temperature storage and thermization on the quality of raw and heat treated milk. National Academic Award 39: 1585 C.F. Dairy Sci., Abet. 55: 2467

دراسة على تحسين بعض صفات الجن الراس المصنوع من لبن الماعز

محاوله للتغلب على رائحة لين الماعز والتي تظهر عادة عند تصنيعه الى جبن جاف (جبن راس). وقد تم في هذه الدراسة تصنيع جبن راس من لين بقرى كفارنه (كتنول) وجبن راس من لين الماعز تم تقسيمه كالآتي:

- (١) لين ماعز معامل حراريا على ٧٥ ° لمدة ١٥ ثانية.

(٢) لين ماعز معامل حراريا على ٨٠ ° لمدة ١٠ ثوانى.

(٣) لين ماعز معامل حراريا على ٨٠ ° لمدة ١٠ ثوانى ثم تبریده الى ٧٥ ° لمدة ٢٤ ساعه.

(٤) لين ماعز معامل حراريا على ٨٠ ° لمدة ١٠ ثوانى ثم تبریده الى ٧٥ ° لمدة ٤٨ ساعه

وتم تخزين جميع الجبن الناتج للتسوية على درجة حراره ١٣° ± ١° ورطوبه نسييه من ٩٠-٨٠ %

لمدة ثلاث شهور تم خلالها تحليل الجبن كيميائيا مع تقدير بعض المجاميع الميكروبيه كما تم تقييمه حسيا كل شهر ولمدة ثلاث شهور كما تم تقدير النسبة المئويه للتصاصي قبل وبعد التقليل وكذلك في نهاية مدة التسوية ويمكن تخيص اهم النتائج المتحصل عليها فيما يلى:

 - (١) زيادة النسبة المئويه للتصاصي في الجبن الناتج من اللين الماعز عن مثيلاتها الناتجه من اللين البقرى.
 - (٢) زيادة نسبة الرطوبه و الدهن و النيتروجين الكلى في الجبن الناتج من اللين الماعز عن الجبن الناتج من اللين البقرى سواء في الجبن الطازج أو خلال مدة التسوية .
 - (٣) زيادة نسبة دلائل التسوية النسبة المئويه للنيتروجين الذائب وكذلك نسبته الى النيتروجين الكلى في جميع الجبن الناتجه من معاملات اللين الماعز عن مثيلاتها الناتجه من اللين البقرى.
 - (٤) لوحظ أيضاً زيادة نسبة الأحماض الدهنية الطياره وكذلك الأحماض الدهنية قصيرة السلسله والتي يعزى إليها أساس الطعم الناضج الجيد في جميع معاملات الجبن الناتج من اللين الماعز عن الجبن الناتج من اللين البقرى.
 - (٥) كما تشير النتائج الى الانخفاض التدريجي في العدد الكلى للبكتيريا وعدد البكتيريا المحطة للدهن وكذلك عدد البكتيريا المحطة للبروتين طول فترات التسوية في جميع المعاملات.

(١) وعند تحكيم الجبن الناتج حسيا حصل الجبن الناتج من اللين الماعز والمعامل حراريا على درجة حراره ٨٠ ° لمدة ١٠ ثوانى وبدون تخزين اللين إلى للتبريد على درجة تحكيم حسيه تقارب درجة الجبن الناتج من اللين البقرى حيث تميز الجبن الناتج من هذه المعامله بالنظافه ولم تلاحظ به رائحة اللين الماعز والتي ظهرت عادة عند تصنيع اللين الماعز الى منتجات لبنيه أخرى كذلك تميز جبن هذه المعامله بارتفاع معدلات التسوية.

تحكيم البحث

أ.د / طه عبد الحليم نصري

أ.د / منير محمود العبد

كلية الزراعة - جامعة المنصورة

كلية الزراعة - جامعة القاهرة