

38. Five weeks after treatment of dendritic corneal ulcer there is appearance of an ulcer with circular raised everted edges, it is:
- a) Metaherpetic ulcer
  - b) Atheromatous ulcer
  - c) Neurotrophic ulcer
  - d) Neuroparalytic ulcer
39. Vision loss in Riley-Day syndrome is often due to:
- a) Cataracts
  - b) Optic nerve hypoplasia
  - c) Amblyopia
  - d) Corneal scarring
40. Which one of the following is associated with cystoid macular edema?
- a) Ecothiopate
  - b) Timolol
  - c) Epinephrine
  - d) Pilocarpine
41. Which concerning xanthelasma is true:
- a) Present most often at lateral aspect of upper lid
  - b) Most affected do not have hyperlipidemia
  - c) The deposits are in deep layers of the dermis
  - d) The deposits are frequently around arterioles
42. Follicular conjunctivitis is more related to which of the following:
- a) Propaine
  - b) Pilocarpine
  - c) Timolol
  - d) Betaxolol
  - e) Carbacol
43. The condition in which same size donor button is used in penetrating keratoplasty:
- a) Bullous keratopathy after cataract surgery
  - b) Keratoconus
  - c) Corneal stromal dystrophies
  - d) Fungal keratitis
44. Antinuclear antibody is helpful in diagnosis of:
- a) Rheumatoid artheritis
  - b) Behcet disease
  - c) Juvenile artheritis
  - d) Ankylosing spondylitis