



On The Formulation of Space-Time Fractional Reaction-Convection-Diffusion Equation in Comb-Like Model

Israa M. Abdelsabour¹, Mohammed Sallah^{1,2}, Mohsen A. Zahran¹, Ahmed Elgarayhi¹

¹ *Physics Department, Faculty of Science, Mansoura University, Mansoura 35516, Egypt*

² *Higher Institute for Engineering and Technology, New Damietta, Egypt*

Received: 13/3/2021
Accepted: 1/4/2021

Abstract: The general equation of fractional reaction-diffusion windswept to an external force expressed as a velocity term suggested to explain the mechanism of transportation in complex systems with time-space variance. This work uses the Caputo derivation form to derive and solve the generalized fractional reaction-diffusion equation in Mittag-Leffler and H-Fox functions. The solution used to present some explanation of the anomalous diffusion phenomena in some disordered systems. For example, there is a sturdy index to show that for a system with doubled trappings, fractional calculus provides some physical view of fractional dynamics like diffusion in the dendritic spine of the nervous system as an example of the diffusion process in biological systems.

keywords: Anomalous diffusion; Comb-like model; Convection term; Fractional reaction-diffusion equation

1. Introduction

Diffusion is a passive transport process that results in a net movement of particles without requiring bulk motion, from higher to lower concentrations. It exists in all states of matter on various time and space scales, which characterizes many physical, chemical, and biological phenomena, based on Fick's laws, or by taking into account a microscopic point of view in terms of the erratic motion of the suspended particles [1]. Diffusion can be described in terms of the mean-squared displacement (MSD) of the particle spreading $\langle x^2 \rangle \propto t$, where x is the distance a particle moved in time t , which is known as normal diffusion (Brownian or Fickian). The standard normal diffusion described by the Gaussian distribution can be obtained from the usual Fokker-Planck equation with a constant diffusion coefficient and zero drift [2].

However, the disparity between anomalous diffusion and normal diffusion in terms of variance indicates a nonlinear increase in time

$$\langle x^2 \rangle \sim t^\gamma \tag{1}$$

For this case, the MSD can be classified via anomalous diffusion exponent; the exponent γ can be either subdiffusion for $0 < \gamma < 1$ or super diffusion for $\gamma > 1$. In recent decades, the anomalous diffusion has been extensively studied in a variety of physical applications, e.g., the fluids transport in porous media [3], surface growth [4], numerical solution of the time-fractional reaction-advection diffusion equation in porous media [5], and the biological population's spatial diffusion [6].

A lot of examples can be found in Refs. [7,8]. To study the anomalous diffusion applications, the conventional Fokker-Planck equation has been used. The standard Fokker-Planck equation, for example, may result in anomalous diffusion but with a variable diffusion coefficient. [9,10]. It is also achieved by subsuming nonlinear terms in the diffusion section or external forces [11,12,13]. However, several approaches to studying anomalous diffusion and the associated phenomena have focused on fractional equations [14,15]. Such interest in the absence and existence of external velocity or force fields has been extensively shown to be a compatible instrument for

explaining anomalous transport processes [8,16,17].

Recently, the anomalous transport phenomenon on living biological cells have been revealed on the mesoscopic scale dynamics. Following the recent high-detailed simulation, this abnormal behavior found in living cells and the environments of artificially crowded regulation operates in the form described in Ref. [7,18].

Moreover, there is a swelling interest in studying the anomalous diffusion equation using fractional calculus, e.g., Elwakil et al. [19] have given a derivation for the fractional space-time diffusion equation using the comb-like structure as a draft model obtaining the solution for three different interesting cases. Zahran [20] described the equation with an integrodifferential form, defining the fractional reaction-diffusion equation's derivation in the general form concerning time and space operators instead of the first-order derivative the Riemann-Liouville derivative of fractional orders shall contain. Schot et al. [21] found the solutions of a generalized diffusion equation that contains space-time fractional derivatives by taking an absorbent (or source) term and a linear external force into account. They were expressed in terms of the Fox H-function, and for some special cases, they are directly related to the Lévy distributions.

As a matter of fact, the ad-hoc representation of fractional operators should be carried with care. Therefore, this phenomenological approach of this type of differential equations should be solidly justified. Hence, this work suppresses the derivation and solution of the generalized fractional reaction convection diffusion equation using Caputo time-derivative of order γ and space fractional derivative of order α in a comb-like model. We have applied the co-moving technique that has yielded the general solution in the form of Mittag-Leffler and H-Fox Function.

2. Fractional Reaction-Convection-Diffusion Equation

The comb model consists of a backbone along the horizontal axis and fingers or teeth along the perpendicular direction [22] (see Fig. (1) for a two-sided comb) [23]. To explain anomalous transport in percolating clusters, a

comb model was implemented. [24,25] It was considered a toy model for a porous medium used for low-dimensional percolation cluster exploration. [22,24]. The comb-like model has been used many times to derive well-known fractional differential equations, as fractional diffusion equation with or without external force. A matrix of diffusion coefficients describes the comb's geometry, which means that the displacement in the x-direction is possible only along the structure axis, i.e., the diffusion happened at [22]. Hence, the main motive of this work is assuming the anomalous diffusion along with the tooth of the structure. One can obtain the generalized form of the space-time fractional reaction-convection-diffusion equation for the sake of completeness. This model can describe the transfer of the nerve signals in the dendritic spines, where the signal particles may be trapped. The fingers of the comb-like model cause the trapping effect. Consequently, the diffusion in the y-direction shall be considered in the convection term as it should be added to the x-direction diffusion




Fig.1. Two-sided comb-like model.

In terms of the probability density function $P(x, y, t)$, let the x-component of the current J along the backbone is given by

$$J_x = \delta(y) \left[v(x)P(x, y, t) - D_x \frac{\partial^\alpha}{\partial x^\alpha} P(x, y, t) \right] \quad (2)$$

Moreover, the corresponding y-component of the current, along with fingers (where the trapping may occur), has the form

$$J_y = -D(t) \frac{\partial}{\partial y} P(x, y, t) \quad (3)$$

The diffusion coefficient $D(t)$ is believed to be a time-dependent quantity. From the continuity equation that follows

$$\frac{\partial}{\partial t} P(x, y, t) + \nabla \cdot J = -\delta(y)\psi(x, y, t) \quad (4)$$

Where $\psi(x, y, t)$ corresponds to the reaction term. By using eq. (2),(3)and (4), one shall get

$$v^*(x, \gamma) = \frac{V(x)\Gamma(\gamma)}{\sqrt{\pi D_1}}, \quad \gamma = 1 - \frac{\beta}{2}$$

$$\left[\frac{\partial}{\partial t} - D(t) \frac{\partial^2}{\partial y^2} \right] P(x, y, t) = \delta(y) \left\{ D_\alpha \frac{\partial^\alpha}{\partial x^\alpha} P(x, y, t) - \frac{\partial}{\partial x} [v(x)P(x, y, t)] - \psi(x, y, t) \right\}$$

$$(5)$$

$$D_\alpha \frac{\partial^\alpha}{\partial x^\alpha} P(x, y, t) - \frac{\partial}{\partial x} [v(x)P(x, y, t)] = \varphi(x, y, t)$$

with $\alpha = \mu + 1$. Assuming (6)

$$\left[\frac{\partial}{\partial t} - D(t) \frac{\partial^2}{\partial y^2} \right] P(x, y, t) = \delta(y) [\varphi(x, y, t) - \psi(x, y, t)] \quad (7)$$

The effective diffusion coefficient is believed to be $D(t)$, that has the explicit time dependency as follows: [26], with the coefficient D_1

$$D(t) = D_1 \beta t^{\beta-1} \quad (8)$$

with the exponent β that represents the diffusion type along with the structure of the comb. $0 < \beta < 1$ Is sub-diffusion that describes a diffusion process slower than normal diffusion. $1 < \beta < 2$ Is super-diffusion, a form of diffusion that is faster than normal diffusion. Diffusion in fingers is a fractional Brownian motion operated by the effective Fokker-Plank equation, and diffusion in the backbone is a time fractal process with a memory kernel. For that reason, the Green function associated with the homogeneous part of the equation. (7) impact by the initial condition $G(y, 0) = \delta(y)$ of has the form

$$G(y, t) = \frac{1}{\sqrt{\pi D_1 t^{\beta-1}}} \exp\left[\frac{-y^2}{4 D_1 t^{\beta-1}} \right]$$

$$s^\gamma P(x, s) - s^{\gamma-1} \delta(x) = D_o \frac{\partial^\alpha}{\partial x^\alpha} P(x, s) - \psi(x, s) \quad (9)$$

The general solution for equation (5) can be defined through the integration over the source term and for the different parameter values $t = t - t'$ as

$$P(x, 0, t) = \int G(y - y', t - t') \left\{ \delta(y') \left[D_\alpha \frac{\partial^\alpha}{\partial x^\alpha} P(x, y', t') - \frac{\partial}{\partial x} [v(x)P(x, y', t')] - \psi(x, y', t') \right] \right\} dy' dt' \quad (10)$$

incorporating equation (9) into equation before integration over y (i.e., along with the principal channel), the closed equation for the concentration of the particles along the main channel [27] is given by

$$P(x, t) = {}_0 D_t^\gamma \left\{ D_\alpha \frac{\partial^\alpha}{\partial x^\alpha} P(x, t) - \frac{\partial}{\partial x} [v^*(x, \gamma)P(x, t)] - \psi(x, t) \right\} \quad (11)$$

under the following circumstance in terms of Gamma function

$$\Gamma(\gamma) \quad (12)$$

where the Caputo derivative ${}_0 D_t^\gamma$ is defined as

$${}_0 D_t^\gamma F(t) = \frac{1}{\Gamma(n - \alpha)} \int_0^t (t - t')^{n - \alpha - 1} F(t') dt'$$

$$H_{p,q}^{m,n} \left[z^\alpha \begin{matrix} (a_p, A_p) \\ (b_q, B_q) \end{matrix} \right] = \frac{1}{\alpha} H_{p,q}^{m,n} \left[z \begin{matrix} (a_p, A_p / \alpha) \\ (b_q, B_q / \alpha) \end{matrix} \right] \quad (13)$$

It is clear here that due to the random walk particle trapping along the y-direction, the fractional-order integration occurs. When the fractional derivative of Caputo is applied on both sides, it produces

$$\frac{\partial^\gamma}{\partial t^\gamma} P(x, t) = D_\alpha \frac{\partial^\alpha}{\partial x^\alpha} P(x, t) - \frac{\partial}{\partial x} [v(x)P(x, t)] - \psi(x, t) \quad (14)$$

3. Analytical Solution of The Fractional Diffusion Problem

Taking the special case that $D_\alpha = D_o$ and $V(x) = V_o$, the fractional reaction advection-diffusion equation can be studied if we convert the transport analysis from an external observer reference frame to a co-moving frame, as

At the origin $t = 0$, we consider a point source so that the initial condition is given by $P(x, 0) = \delta(x)$. Taking the Laplace transform of equation (15) gives

$$\text{We suppress the imaginary unit in Fourier space [8] as}$$

$$F \left\{ {}_0 D_x^\alpha F(x) \right\} = -|k|^\alpha F(x) \quad (16)$$

(17)

Then, equation (16) will have the form

$$p(k, s) = \frac{s^{\gamma-1}}{s^\gamma + D_0 |k|^\alpha} - \frac{\psi(k, s)}{s^\gamma + D_0 |k|^\alpha} \quad (18)$$

Taking the inverse Laplace and inverse Fourier to the above equation, respectively, will give the whole solution. Accordingly, the general solution of the probability density function [28] can be expressed as

$$P(x, t) = \int_{-\infty}^{\infty} G_1(x-x', t) dx' + \int_0^t (t-t')^{\beta-1} \int_0^x \psi(x', t') G_2(x-x', t-t') dx' dt' \quad (19)$$

where,

$$G_1(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp(-ikx) E_{\gamma,1}(-D_0 |k|^\alpha t^\gamma) dk \quad (20)$$

$$E_{\alpha,\beta}(z) = H_{12}^{11} \left[-z \left| \begin{matrix} (0,1) \\ (0,1) \end{matrix} \right. (1-\beta, \alpha) \right]$$

(21)

where H-function is defined by

(22)

Taking the inverse cosine Fourier transform of the Fox H-function in the form [28]

(23)

Consequently, in terms of the Fox H-function, we get the analytical solution of the

$$\int_0^{\infty} t^{\rho-1} \text{Cos}(kt) H_{p,q}^{m,n} \left[a t^\mu \left| \begin{matrix} (a_p, A_p) \\ (b_q, B_q) \end{matrix} \right. \right] dt = \frac{\pi}{k^\rho} H_{q+1, p+2}^{n+1, m} \left[\frac{k^\mu}{a} \left| \begin{matrix} (1-b_p, B_p) & \left(\frac{1+\rho}{2}, \frac{\mu}{2} \right) \\ (\rho, \mu) & (1-a_p, A_p) & \left(\frac{1+\rho}{2}, \frac{\mu}{2} \right) \end{matrix} \right. \right]$$

space-time fractional advection-reaction diffusion equation, as

$$G_1(x, t) = \frac{1}{|x-vt|^\alpha} H_{13}^{2,1} \left[\frac{|x-vt|}{D_0^{1/\alpha} t^{\gamma/\alpha}} \left| \begin{matrix} (1,1/\alpha) & (1, \gamma/\alpha) & (1,1/2) \\ (1,1) & (1,1/\alpha) & (1,1/2) \end{matrix} \right. \right] \quad (24)$$

and

$$G_2(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp(-ikx) E_{\gamma,\gamma}(-D_0 |k|^\alpha t^\gamma) dk \quad (25)$$

whose form can express in terms of Fox H-function, as

$$G_2(x, t) = \frac{1}{|x-vt|^\alpha} H_{13}^{2,1} \left[\frac{|x-vt|}{D_0^{1/\alpha} t^{\gamma/\alpha}} \left| \begin{matrix} (1,1/\alpha) & (\gamma, \gamma/\alpha) & (1,1/2) \\ (1,1) & (1,1/\alpha) & (1,1/2) \end{matrix} \right. \right] \quad (26)$$

$G_1(x, t)$ and $G_2(x, t)$ Can give the general

solution of the probability density function in terms of H-Fox Function.

4. Conclusion

We have derived and analyzed the fractional reaction convection diffusion equation with Caputo time-derivative of the order and space fractional derivative of the order in a comb-like model. The random walker diffuses along the x-axis, and on the y-axis, the trapping events occur. The diffusion along the x-direction is a process of fractal time that showed a mean-squared displacement $\langle x^2 \rangle \propto t^\gamma$ with Delta function's initial condition $P(x, 0) = \delta(x)$. Moreover, the y-direction diffusion is given by mean-squared displacement $\langle y^2 \rangle \propto t^\beta$, with order β related to order γ (see equation (12)). The main point here is the direct link between the integrated fractional operators and the random disappearances or trappings along with the structural fingers. Also, we have introduced the co-moving technique in which the Laplace-Fourier technique can be applied to the resulting equation to give the solution in the form of Fox H-Function to get the solution described by the equation (19) – equation (26). The studied model can describe some biophysical systems with trapping phenomena like the diffusion in the dendritic spine of neurons.

4. References

- 1 Bressloff P. C., J. M. Newby, (2013) Stochastic models of intracellular transport, *Rev. Mod. Phys*, **85**(1), , 135-196.
- 2 Risken H., (1984) *the Fokker-Planck equation* (Springer, **18**, , 63-95).
- 3 Muskat M., (1946). the flow of homogeneous fluids through porous media (McGraw-Hill Book Company, New York).
- 4 Spohn H. , (1993), Surface dynamics below the roughening transition, *J. Phys. I France* **3**(1), 69-81.
- 5 Pandey P., S. Kumar, J.F. Gomez-Aguilar, (2020) Numerical Solution of the Time Fractional Reaction-advection diffusion Equation in Porous Media, *J. Appl and comput mech*, **12**(10),.
- 6 Morton E., C. MacCamy, (1977), On the diffusion of biological populations,

- 7 *Mathematical Biosciences*, **33(1-2)**, 35-49.
- 8 Bouchaud J. P., A. Georges, Anomalous diffusion in disordered media: Statistical mechanisms, models and physical applications, *Phys Rep*, **195(4-5)**, 1990, 127-293.
- 9 Metzler R., J. Klafter, (2000)The random walk's guide to anomalous diffusion: A fractional dynamics approach, *Phys Rep*, **339(1)**, 1-77.
- 10 Fa K. S., (2005). Exact solution of the Fokker-Planck equation for a broad class of diffusion coefficients, *Phys. Rev. E* **72(2)**,
- 11 Fa K. S.,(2011)Solution of Fokker-Planck equation for a broad class of drift and diffusion coefficients, *Phys. Rev. E* **84(1)**,
- 12 Bologna M., C. Tsallis, and P. Grigolini, (2000). Anomalous diffusion associated with nonlinear fractional derivative Fokker-Planck-like equation: Exact time-dependent solutions, *Physical Review E* **62(2)**,
- 13 Assis Jr., P. da Silva, L. da Silva, E. Lenzi, and M. Lenzi, (2006) *Journal of mathematical physics* **47(10)**,
- 14 Lenzi E., M. Lenzi, T. Gimenez, and L. da Silva, (2010), *Journal of Engineering Mathematics*, **67(3)**, 233-240.
- 15 Sokolov I. M., J., Klafter, and A. Blumen, (2002), Fractional Kinetics, *Physics Today*, **55(11)**, 48-54.
- 16 Tsallis C., and E. Lenzi, Anomalous (2002), diffusion: Nonlinear fractional Fokker-Planck equation, *Chemical Physics*, **284(1-2)**, 341-347.
- 17 El-Wakil S. A., M. A. Zahran, (2001)Fractional representation of Fokker-Planck equation, *Chaos, Solitons and Fractals*, **12(6)**, 1929-1935.
- 18 El-Wakil S. A., M. A. Zahran, M. A. Abdou, (2003)The operator method for solving the fractional Fokker-Planck equation, *J Quant Spectrosc Radiat Transf*, **77(3)**, 317-327.
- 19 Havlin S., D. Ben-Avraham, (1987), Diffusion in disordered media, *Adv Phys*, **36(6)**, 695-798.
- 20 El-Wakil S. A., M. A. Zahran, E. Abuelwafa, (2004), Fractional (space-time) diffusion equation on comb-like model, *Chaos, Solitons and Fractals*, **20(5)**, 1113-1120.
- 21 Zahran M. A., (2009) On the derivation of fractional diffusion equation with an absorbent term and a linear external force, *Applied Mathematical Modelling*, **33(7)**, 3088-3092.
- 22 Schot A., M. K. Lenzi, L. R. Evangelista, L. C. Malacarne, R. S. Mendes, E. K. Lenzi,(2007), Fractional diffusion equation with an absorbent term and a linear external force: Exact solution, *Physics Letters A*, **366(4-5)**, 346-350.
- 23 Arkhincheev V., E. Baskine, A. Tybulewicz, Anomalous diffusion and drift in a comb model of percolation clusters, *Sov. Phys. JETP*, **73(1)**, 1991, 161-165.
- 24 Sandev T., A. Iomin, V. Méndez,(2016)Lévy processes on a generalized fractal comb. *J Phys A Math Theor*, **49(35)**,
- 25 White S. R., M. Barmat,(1984)Field-induced drift and trapping in percolation networks, *J. Phys. A: Math. Gen*,**17(15)**,
- 26 Weis G. H., S. Havlin, (1986), Some properties of a random walk on a comb structure, *Phys A Stat Mech its Appl*, **134(2)**, 474-482.
- 27 Wang K. G., C. W. Lung,(1990), Long-time correlation effects and fractal Brownian motion, *Phys. Lett. A*, **151(3-4)**, 119-121.
- 28 Arkhincheev V., Anomalous diffusion and charge relaxation on comb model: exact solutions, *Physica A*,**280(3)**, 304-314.
- 29 Haubold H. J., A. M. Mathai, R. K. Saxena, (2007), Solution of fractional-reaction diffusion equation in terms of H-function, *math. PR*, **35**, 681-689.