

GENETICAL ANALYSIS FOR F₁ AND F₂ GENERATION IN SOME EGYPTIAN COTTON CROSSES (*Gossypium barbadense* L.)

Nazmey, M.N.A.; A.E.M. Eissa and W.M.B. Yehia

Cotton Research Institute, Agriculture Research Center. Egypt

ABSTRACT

Five cotton varieties belong to *Gossypium barbadense* L. i.e. Giza 89, Giza 90, Giza 83, Pima S₄ and Pima S₆ were selected as parents and crossed in a half diallel pattern to evaluate general and specific combining ability effects (GCA and SCA) and heterotic effects for some agronomic traits i.e. boll weight (B.W.), seed cotton yield (S.C.Y.), lint yield (L.C.Y.), lint percentage (L.%), seed index (SI), length at 2.5% , strength g/tex and micronaire value (Mic). Analysis of variance revealed significant differences among entries for all traits studied except for B.W., S.I. and Mic in F₁ and F₂ generations as well as length at 2.5% in F₂'s generation which showed significant differences. The mean squares in both F₁'s and F₂'s for general combining ability (GCA) were insignificant for all traits except for S.C.Y. and L.C.Y. in F₁'s and F₂'s generations. Meanwhile, the mean squares for specific combining ability (SCA) were insignificant for all traits except S.C.Y. and L.C.Y. in both generations and strength g/tex in F₁'s generation which showed significant. The GCA/SCA ratio of variance components indicated that additive genetic variance was generally importance for B.W., L%, SI and Mic in the F₁ hybrids and for B.W., L.C.Y., L.% and Mic in the F₂ generation. Mid-parents heterosis values were significant and positive for S.C.Y., L.C.Y. and length at 2.5% in the cross Giza 89 x Giza 90 (P₁ x P₂), S.C.Y., L.C.Y., L% and length at 2.5% in the cross Giza 89 x Giza 83 (P₁ x P₃), S.C.Y., S.I., length at 2.5% and Mic in the cross Giza 89 x Pima S₄ (P₁ x P₄), SCY, L.C.Y. and length at 2.5% cross Giza 89 x Pima S₆ (P₁ x P₅), SCY, LY and Mic in the cross Giza 90 x Giza 83 (P₂ x P₃), S.C.Y., L.C.Y. and length at 2.5% in the cross Giza 90 x Pima S₄ (P₂ x P₄), length and Mic in the cross Giza 83 x Pima S₄ (P₃ x P₄) and S.C.Y. in the cross Pima S₄ x Pima S₆ (P₄ x P₅). On the other hand, significant negative heterotic values were observed for B.W. and S.I. in the cross Giza 89 x Giza 90 (P₁ x P₂), B.W. in the crosses of Giza 89 x Pima S₄ (P₁ x P₄) and Giza 90 x Giza 83 (P₂ x P₃). Better-parents heterosis values were significant and positive for 1% in the cross Giza 89 x Giza 83 (P₁ x P₃), S.I. and length cross Giza 89 x Pima S₄ (P₁ x P₄), S.C.Y. in the cross Giza 89 x Pima S₆ (P₁ x P₅), L.C.Y. in the cross Giza 90 x Giza 83 (P₂ x P₃) and S.C.Y. and L.Y. in the cross Giza 90 x Pima S₄ (P₂ x P₄). These results indicated to the importance of specific combining ability in the genetic expression of these traits with respect to the studied crosses. Regarding inbreeding depression, significant positive effect were obtained for L.C.Y. in the cross (P₁ x P₂), B.W., S.C.Y. and L.C.Y. in the cross (P₁ x P₃), S.C.Y. and S.I. in the cross (P₁ x P₄), S.C.Y., L.Y. and L.% in the cross (P₁ x P₅), S.C.Y. and L.C.Y. in the cross (P₂ x P₃), S.C.Y., L.C.Y. and L.% in the cross (P₂ x P₄), S.C.Y. in the cross (P₂ x P₅) and in the cross (P₃ x P₄) and S.C.Y. in the cross (P₄ x P₅). This finding revealed the importance of heterotic effect in these traits with respect to the studied crosses .

INTRODUCTION

Diallel analysis is one of the methods that reveal yield potentiality of the cotton cultivars and their crosses on the basis of their general and specific combining ability. General combining ability (GCA) includes the additive variance, while specific combining ability (SCA) could be considered

as a measure of non-additive genetic variance arising largely from dominance and epistatic deviations. Several studies have been established in this respect by many investigators.

El-Dobaby *et al.* (1997) found highly significant effect of GCA and SCA for each of seed cotton yield/plant, boll weight, lint percentage, seed index and lint index. Hendawy *et al.* (1999) reported that both GCA and SCA were highly significant in all studied fiber attributes.

Khorgade *et al.* (2000) and Zia *et al.* (2001) determined GCA and SCA in seven American cotton genotypes, they indicated that the GCA and SCA were highly significant for ginning percentage, lint index, seed index, micronaire value and upper half means. El-Adl *et al.* (2001) revealed that GCA were highly significant for boll weight and ginning out turn, while SCA were highly significant for yield and staple length. Laxman and Genesh (2003) revealed that SCA variance were higher for boll weight, seed cotton yield, seed and lint index and halo length than GCA. Esmail *et al.* (2005) studied combining ability in some Egyptian cotton genotypes, they found that significant positive GCA effects with regard to seed cotton yield and most of its contributing variables. El-Adly (2008) found highly significant effects for GCA for seed cotton yield/plant, lint yield, lint percentage, seed index and upper half mean, while he found highly significant for SCA for lint percentage, seed index, lint index and upper half mean.

The objective of this investigation is to study the relative magnitude of additive and non-additive genetic variance through evaluation both general and specific combining ability (GCA and SCA) effects for yield and yield components in diallel crosses.

MATERIALS AND METHODS

A half diallel of five cotton cultivars namely, Giza 89, Giza 90, Giza 83, Pima S₄ and Pima S₆ belong to *G. barbadense* L. were evaluated for seed cotton yield and some agronomic characters. In 2007 season, the five parents were grown and all possible crosses according to half diallel mating design were carried out. In 2008 season, the 10 F₁'s hybrid seeds were planted in order to obtain the F₂'s generation through self-fertilization. The parental varieties were also crossed to obtain additional F₁'s hybrid seeds. The F₁'s seeds and F₂'s seeds were produced at Seds Experimental Station, Agricultural Research Center at Bany Souif governorate. In 2009 season, a randomize complete blocks trial with three replicates was carried out including the five parental varieties and ten F₁'s and F₂'s populations in Seds Experimental Station. Each plot was two rows 7 m long and 60 cm apart, the space between hills 50 cm. The hills were thinned to one plant/hill. Cultural practices were carried out as usually done in Seds Experimental Farm. Eight characters were studied, i.e.

1. Boll weight (BW), average weight in grams.
2. Seed cotton yield/plant (SCY) in grams.
3. Lint cotton yield/plant (LY) in grams.

4. Lint percentage (L%). $L\% = \frac{\text{Lint yield}}{\text{Seed cotton yield}} \times 100$
5. Seed index (SI) in grams.
6. Length in mm
7. Strength (G/tex)
8. Micronaire reading (Mic)

Estimates of combining ability were carried out according to Griffing's (1956) method 2 model 1 and were analyzed on a plot mean basis to obtain estimates of general and specific combining ability (GCA and SCA) effects and variances. All effects were assumed to be fixed.

Heterosis was expressed for all studied traits as percent increase of the F₁'s performance above the mid-parents (M.P.) and better parents (B.P.) values. Inbreeding depression was calculated from comparison between F₁ and F₂ generations.

RESULTS AND DISCUSSION

Analysis of variance:

The analysis of variance for genotypes, general combining ability (GCA) and specific combining ability (SCA) in addition to GCA/SCA ratio are presented in Table 1 for two populations (F₁ and F₂). The results showed that the differences among genotype were significant or highly significant for all traits in both populations (F₁'s and F₂'s) except for boll weight (B.W.), seed index (S.I.) and micronaire values (Mic) in F₁'s and F₂'s. Mean squares of GCA and SCA showed that the GCA were highly significant for seed cotton yield (S.C.Y.) and lint yield (L.C.Y.) for the two populations (F₁'s and F₂'s). The mean squares of SCA were significant or highly significant for seed cotton yield (S.C.Y.), lint yield (L.C.Y.) and the strength G/tex in F₁'s and lint yield (L.C.Y.) and strength G/tex in F₂'s.

Table 1: Mean squares for genotypes and combining ability (GCA and SCA) in F₁'s and F₂'s generations for studied traits.

S.O.V.		B.W.	S.C.Y.	L.C.Y.	L.%	S.I.	Length	Strength G/tex	Mic
Genotypes	F ₁	0.095	762.811**	142.759**	2.918*	0.447	3.571**	10.517**	0.550
	F ₂	0.201	1304.6**	110.831**	2.754**	0.407	2.003	7.544**	0.389
General combining ability (GCA)	F ₁	0.059	1017.213**	20.756**	1.514	0.222	1.058	3.057	0.380
	F ₂	0.081	286.441**	81.860**	2.089	0.104	0.499	1.780	0.168
Specific combining ability (SCA)	F ₁	0.021	313.093**	58.318**	0.756	0.120	1.243	3.685*	0.105
	F ₂	0.061	494.251	18.977**	0.916	0.149	0.735	2.809*	0.114
GCA/SCA	F ₁	2.810	0.342	0.356	2.003	1.850	0.851	0.830	3.619
	F ₂	1.328	0.580	4.314	2.281	0.698	0.679	0.634	1.474

*,** significant at 0.05 and 0.01 levels of probability, respectively

GCA/SCA ratio indicated that the GCA was greater than SCA for all studied traits in the two generations except for seed cotton yield (S.C.Y.), lint yield (L.C.Y.), length mm and strength G/tex in F₁'s and seed cotton yield (S.C.Y.), seed index (S.I.), length and strength G/tex in F₂'s generations. Therefore, it could be concluded that most of the genetic variance for those traits was due to additive and non-additive gene actions. These results are in

general agreement with those reported by Rahoumah and El-Shaarawy (1992), Khorgade *et al.* (2000), El-Adl *et al.* (2001), Laxman and Genesh (2003) and Ismail *et al.* (2005).

Combining ability:

General combining ability effects (GCA) of the parents for each trait are presented in Table 2. The results showed that the parent Giza 89 showed significant and positive GCA effects for boll weight (B.W.) and length in F₁'s. While, it was highly significant and positive GCA effects for seed index (S.I.) and micronaire values (Mic) in F₁'s and for boll weight (B.W.) and micronaire value (Mic) in F₂'s. So GCA effects were negative and significant for seed cotton yield (in F₁'s and F₂'s), lint yield (L.C.Y.), lint percentage (L%) in F₁'s, while it was negative and highly significant for lint yield (L.C.Y.) and lint percentage (L%) in F₂'s.

Table 2: Parental mean performances and mean estimates of GCA effects of five parents and their F₁'s and F₂'s generations.

Traits	Generations	P ₁ (G89)	P ₂ (G90)	P ₃ (G83)	P ₄ (PS ₄)	P ₅ (PS ₆)	L.S.D.		g [^] _i	g [^] _i r [^] g [^] _{ij}
							1%	5%		
B.W.	\bar{X}	3.1	3.1	2.8	2.7	2.7				
	F ₁	0.089*	0.98**	-0.007	-0.088*	-0.092**	0.091	0.067	0.033	0.420
	F ₂	0.101**	0.101**	-0.142*	0.015	-0.075*	0.099	0.073	0.036	0.458
S.C.Y.	\bar{X}	135.5	155.6	177.3	161.7	169.6				
	F ₁	-6.043*	-0.905	4.557	0.805	1.586	7.245	5.370	2.622	33.511
	F ₂	-10.446*	1.778	6.135	-1.031	3.564	13.721	10.170	4.966	63.465
L.C.Y.	\bar{X}	48.5	61.9	65.5	61.2	63.8				
	F ₁	-2.777*	1.099	1.704	-0.149	0.123	2.880	2.134	1.042	13.319
	F ₂	-5.307**	0.489	2.827*	-1.054	3.046*	3.068	2.274	1.110	14.192
L.%	\bar{X}	35.8	39.8	36.9	37.8	37.6				
	F ₁	-0.351*	0.815**	-0.099	-0.166	-0.199	0.446	0.331	0.162	2.064
	F ₂	-0.683**	0.731**	0.155	-0.373*	0.170	0.416	0.308	0.151	1.924
S.I.	\bar{X}	9.8	9.9	9.3	9.5	9.0				
	F ₁	0.254**	0.088	-0.089	-0.041	-0.212**	0.169	0.125	0.061	0.782
	F ₂	-0.010	0.205**	-0.062	-0.019	-0.114	0.178	0.132	0.065	0.118
Length	\bar{X}	31.0	31.0	30.8	31.0	31.7				
	F ₁	0.520*	-0.318	-0.309	0.310	-0.204	0.536	0.398	0.194	2.481
	F ₂	0.036	-0.297	-0.083	-0.083	0.427	0.602	0.446	0.218	2.782
Strength G/tesx	\bar{X}	33.6	37.6	38.8	37.6	35.9				
	F ₁	-0.970	-0.265	0.478	0.711	0.045	1.485	1.100	0.537	6.867
	F ₂	-0.631	-0.408	0.583	0.292	0.164	1.315	0.975	0.476	13.519
Mic	\bar{X}	4.1	4.1	3.3	3.1	4.2				
	F ₁	0.293**	0.155*	-0.164**	-0.283**	-0.002	0.164	0.122	0.059	0.759
	F ₂	0.176**	0.062	-0.071	-0.229**	0.062	0.156	0.115	0.056	0.721

*,**significant at 0.05 and 0.01 levels of probability, respectively .

Meanwhile, the parent Giza 90 showed significant, highly significant and positive GCA effects for boll weight (B.W.), lint percentage (L.%) in both generations, seed index (S.I.) in F₂'s and micronaire value in F₁'s. Parent Giza 83 showed insignificant GCA effects for all traits in both generations except for boll weight (B.W.) which showed highly significant negative GCA in

F₂'s, lint yield (L.C.Y.) which showed significant positive GCA in F₂'s and micronaire values which showed significant negative GCA in F₁'s. Meanwhile, parent Pima S₄ (P_{S4}) showed insignificant GCA effects for all traits except boll weight (B.W.) which showed significant negative GCA in F₁'s, lint percentage (L.%) which showed significant negative GCA in F₂'s and micronaire values which showed highly significant negative GCA in both generations. Parent Pima S₆ (P_{S6}) showed insignificant GCA effects for all traits except boll weight (BW) which showed highly significant and significant negative GCA in F₁'s and F₂'s, lint yield (L.C.Y.) which showed significant positive GCA in F₂'s and seed index (S.I.) which showed highly significant negative GCA in F₁'s.

It is worth noting that, estimates of GCA effects either positive or negative would indicate that a given parent is much better or much poor than the average of the group involved in the diallel system.

Table 3 showed the SCA effects for each of the ten combinations crosses. From those results, it could be noticed that the cross (P₁ x P₂) showed insignificant SCA effect for all traits except boll weight (B.W.) which showed highly significant negative SCA in both generations and seed index (S.I.) which showed highly significant negative in F₁'s generation. The cross (P₁ x P₃) showed highly positive SCA for seed cotton yield (S.C.Y.) in F₁'s, lint yield (L.C.Y.) in both generations, lint percentage in F₁'s, seed index (S.I.) in F₁'s, length in F₁'s and strength g/tex in F₁'s generation, while it showed highly significant negative SCA effects for boll weight (BW) in F₂'s generation. Meanwhile, the crosses (P₁ x P₄) and (P₁ x P₅) showed highly significant positive SCA effects for most of studied traits. While, it showed highly significant negative SCA effects for boll weight (B.W.) in F₁'s, seed cotton yield (S.C.Y.) in F₂'s, lint yield (L.C.Y.) in F₂'s and seed index in F₂'s generation. On the other hand, showed significant negative SCA effects for boll weight (BW) in F₁'s generation while it showed significant positive SCA effect for strength g/tex in F₂'s generation meanwhile, it showed highly significant positive SCA effect for boll weight (B.W.) in F₂'s, seed cotton yield (S.C.Y.) in F₁'s, lint yield (L.C.Y.) in F₁'s, lint percentage (L%) in F₁'s and seed index (S.I.) in F₂'s generation. The cross (P₂ x P₃) showed highly significant negative SCA effect for boll weight (B.W.) and span length at 2.5% (2.5%S.L.) in F₁'s, while, it showed significant negative SCA for seed cotton yield in F₂'s. Meanwhile, it showed highly significant positive SCA for seed cotton yield (SCY) in F₁'s, lint yield (L.C.Y.) in F₁'s, seed index (SI) in F₂'s and micronaire (M.C.) value in F₁'s generation.

The cross (P₂ x P₄) showed significant and highly significant positive SCA for boll weight (BW) in both generations, seed cotton yield (S.C.Y.) in F₁'s, lint yield (L.C.Y.) in F₁'s and length at 2.5% in F₁'s generation, meanwhile, it showed highly significant negative SCA for lint yield (L.C.Y.) in F₂'s, lint percentage (L%) in F₂'s and strength G/tex in F₂'s generation. The cross (P₂ x P₅) exhibited highly significant positive SCA effects for lint yield in F₂'s and seed cotton yield in F₁'s generation, meanwhile, it showed significant and highly significant negative SCA effects for lint percentage in both generations and micronaire value in F₁'s generation.

The cross ($P_3 \times P_4$) showed highly significant positive SCA effects in both generations for boll weight (B.W.), while, it showed highly significant positive SCA for seed cotton yield (S.C.Y.), lint yield (L.C.Y.) and length in F_1 's generation, meanwhile, it showed highly significant negative SCA in F_2 's generation for lint yield (L.C.Y.), lint percentage (L.%) and strength G/tex. The results showed that the cross ($P_3 \times P_5$) was significant and highly significant negative SCA effects for most of studied traits in F_2 's and length in F_1 's generation. The cross ($P_4 \times P_5$) showed highly significant negative SCA for some of studied traits in F_1 's generation, meanwhile, it showed highly significant positive SCA effects for seed index in F_2 's and the strength G/tex in F_1 's.

The cross which showed significant positive SCA effects could be considered promising crosses for improving these traits.

Heterosis and inbreeding depression:

Table 4 revealed mid-parents and better parent heterotic effects for the characters studied. Concerning the cross ($P_1 \times P_2$), negative significant and highly significant heterotic effects relative to mid-parents were found for boll weight (B.W.) and seed index (SI), while it was positive significant and highly significant for seed cotton yield (S.C.Y.), lint yield (L.C.Y.) and length. Meanwhile, heterosis effects to better parent were negative and significant for B.W., L.% and SI. Regarding the cross ($P_1 \times P_3$) positive significant and highly significant heterotic effects relative to mid-parents were found for S.C.Y., L.C.Y., L.% and length and better parents for L.% and length. The cross ($P_1 \times P_4$) showed negative significant and highly significant heterotic effects relative to mid-parent and better parent for boll weight (B.W.), while it showed positive significant and highly significant relative to mid-parent for S.C.Y., S.I., length and micronaire value (Mic). All traits studied in the cross ($P_1 \times P_5$) showed insignificant heterotic effects relative to mid-parents except S.C.Y., L.C.Y. and length which showed positive significant and highly significant. While, heterosis effects to better parent were negative and highly significant for boll weight (B.W.), meanwhile, it showed positive significant for seed cotton yield (S.C.Y.) and lint yield (L.C.Y.) with respect to the cross ($P_2 \times P_3$) (Table 4). Negative significant heterosis effects were detected for BW, while it showed positive significant and highly significant heterosis effects for S.C.Y., L.C.Y. and micronaire value relative to mid-parents, while, it showed negative significant and highly significant for BW and SI. Meanwhile, it showed positive significant for LY relative to better parents. The cross ($P_2 \times P_4$) showed positive and significant for L.C.Y. relative to better parents. The cross ($P_2 \times P_4$) showed positive and significant heterosis effects relative to mid parents for S.C.Y., L.C.Y. and the length, while better parent heterosis effects were positively significant for seed cotton yield and lint yield and negatively significant for lint percentage (L.%). Concerning the cross ($P_2 \times P_5$), insignificant heterotic effects relative to mid and better parents for all traits except lint percentage (L.%) which showed negative and highly significant relative to better parent. The cross ($P_3 \times P_4$) showed insignificant heterotic effects relative to mid and better parents for all traits except length and micronaire value which showed positive and significant heterosis to mid parents. With respect to cross ($P_3 \times P_5$) insignificant heterosis effects were

detected for all traits. The cross ($P_4 \times P_5$) showed insignificant heterosis effects relative to mid parents for all traits except SCY which showed positive and significant, while seed index (S.I.) and micronaire value traits revealed negative significant and highly significant heterosis to better parent (Table 4).

Table 4: Heterosis value (%) over both mid parents (M.P.) and better parent (BP) and inbreeding depression (ID) for studied traits.

Crosses	Parameters	Characters							
		B.W.	S.C.Y.	L.C.Y.	L.%	S.I.	Length	Strength G/tex	Mic.
$P_1 \times P_2$	H.M.P.	-11.23**	19.70**	20.83**	1.28	-5.08*	5.00*	-3.84	8.87
	H.B.P.	-11.70*	11.95	7.70	-3.85*	-5.72*	4.94	-8.95	8.87
	I.D.	3.571	22.560	18.741*	3.133	-3.226	2.761	-6.122	6.667
$P_1 \times P_3$	H.M.P.	-1.48	22.34**	26.63**	3.68*	3.57	8.44**	4.33	6.65
	H.B.P.	-5.32	7.73	13.39	5.33**	2.39	8.16**	-1.29	-0.81
	I.D.	16.667**	32.304**	14.286*	4.370*	5.000	4.762	3.394	-12.195*
$P_1 \times P_4$	H.M.P.	-9.97*	12.91*	10.74	-1.71	7.20**	8.43**	-5.28	21.82**
	H.B.P.	-15.96**	9.98	7.24	-2.38	5.46*	8.16**	-7.09	8.06
	I.D.	-26.923**	44.407**	27.286	2.981	13.592**	1.190	-8.571	8.889
$P_1 \times P_5$	H.M.P.	-7.19	21.86**	24.27**	2.09	1.83	4.93*	-0.02	11.31
	H.B.P.	-14.89**	14.94*	17.25*	2.04	-1.02	3.05	2.32	0.00
	I.D.	-14.815*	40.585**	23.396**	5.208**	-3.093	0.920	-4.632	4.762
$P_2 \times P_3$	H.M.P.	-9.60*	15.21**	18.79**	0.56	-2.78	-1.24	-8.46	14.41*
	H.B.P.	-13.98**	8.16	15.58*	-3.10	-5.72*	-1.51	-9.80	2.42
	I.D.	-3.704	34.411**	20.608**	1.295	-8.60**	-4.262	-4.286	4.762
$P_2 \times P_4$	H.M.P.	1.56	14.51*	14.69*	0.06	-0.93	5.82	-1.96	5.38
	H.B.P.	-6.45*	16.74*	16.42*	-4.02*	-4.38	5.59	-0.97	-10.48
	I.D.	-13.793	33.210**	25.659**	6.806**	-5.263	5.810*	9.920	-5.405
$P_2 \times P_5$	H.M.P.	0.89	7.74	6.30	-1.40	3.27	-1.34	-7.58	4.98
	H.B.P.	-8.60	5.50	5.18	5.78**	-1.68	-3.05	-7.97	-7.94
	I.D.	-3.571	23.197**	-2.086	0.000	2.062	-6.189	-6.358	-2.564
$P_3 \times P_4$	H.M.P.	-1.22	7.09	7.47	0.36	-1.60	4.96	-6.98	17.71*
	H.B.P.	-3.57	2.39	3.97	-0.88	-2.46	4.74	-8.34	15.31
	I.D.	3.704	26.171	9.838	2.667	-5.435	1.852	-6.197	-10.526
$P_3 \times P_5$	H.M.P.	-0.41	-3.15	-4.32	-1.19	0.00	-2.43	-6.7	4.72
	H.B.P.	-3.57	-7.39	-7.23	-1.60	-0.36	-4.00	-8.94	-11.90
	I.D.	3.704	25.518	0.329	-3.514*	0.00	-8.882*	-4.249	-10.811
$P_4 \times P_5$	H.M.P.	-6.25	11.78*	9.58	-1.99	-4.50	3.03	9.39	-10.00
	H.B.P.	-6.25	9.20	7.37	-2.29	-6.69*	1.89	6.82	-21.43**
	I.D.	-8.00	28.024*	6.131	0.811	-13.636	-9.29	3.483	-18.182*

*,** significant at 0.05 and 0.01 levels of probability, respectively .

Inbreeding depression (ID%) effects were calculated for each cross. The cross ($P_1 \times P_2$) showed insignificant inbreeding depression for characters studied except lint yield (L.C.Y.) which showed positive significant inbreeding depression. Regarding the cross ($P_1 \times P_3$) positive significant and highly significant inbreeding depression effects were found for boll weight (B.W), seed cotton yield (S.C.Y.), lint yield (L.C.Y.) and lint percentage (L.%) while it showed negative and significant inbreeding depression for micronaire value (Mic). Cross ($P_1 \times P_4$) showed positive highly significant inbreeding depression for S.C.Y. and S.I., while it showed negative and highly significant for BW. With respect to the cross ($P_1 \times P_5$) insignificant inbreeding depression for SI, length, strength G/tex and Mic, while B.W. recorded negative

significant inbreeding depression effect and S.C.Y., L.C.Y. and L.% showed positive highly significant inbreeding depression. In cross ($P_2 \times P_3$) showed positive highly significant inbreeding depression for SCY and LY, while, it showed negative highly significant inbreeding depression for seed index (SI). In cross ($P_2 \times P_4$) we showed positive significant and highly significant inbreeding depression for seed cotton yield (S.C.Y.), lint yield (L.C.Y.), lint percentage (L.%) and length. With respect to the cross ($P_2 \times P_5$) (Table 4). Insignificant inbreeding depression (ID%) effects for all traits except S.C.Y. which showed positive and highly significant and length which showed negative significant inbreeding depression. Insignificant inbreeding depression showed in cross ($P_3 \times P_4$) for all characters except S.C.Y. which showed positive significant inbreeding depression. Cross ($P_3 \times P_5$) showed insignificant inbreeding depression for all characters except lint percentage (L.%) and length which showed negative significant and highly significant inbreeding depression. Concerning the cross ($P_4 \times P_5$), insignificant inbreeding depression effects relative to all traits except S.C.Y. which showed positive significant inbreeding depression while SI and Mic showed negative significant and high significant inbreeding depression. The above results indicated that insignificant ID% may be due to the presence of linkage between genes in these materials. In general, the present investigation revealed that not only additive but also non-additive genetic variances were important in the inheritance of yield and yield components characters in cotton breeding programs. Therefore, it could be concluded that recurrent selection program is a proper for improvement these traits with respect to the studied characters.

REFERENCES

- El-Adl, A.M.; Z.M. El-Diasty; A.A. Awaad; A.M. Zeina and A.M.A. El-Bary (2001). Inheritance of quantitative traits of Egyptian cotton (*Gossypium barbadense* L.). A yield and yield component traits. Egypt. J. Agric. Res. 79(2): 625-646.
- El-Adly, H.H. (2007). Heterosis, inbreeding depression and combining ability analysis in some cotton crosses (*Gossypium barbadense* L.). Egypt. J. Agric. Res., 86(3): 1081-1093.
- El-Debaby, A.S.; M.M. Kassam; M.M. Awaad and G.M. hemaida (1997). Heterosis and combining ability in intervarietal crosses of Egyptian cotton in different locations. Egypt. J. Agric. Res., 75(3).
- Esmail, E.M.; H. El-Mahfouz and M.D.H. Dewedar (2005). Mean performance, combining ability and heterosis of new Egyptian cotton genotypes as parental genotypes in breeding programs. Egypt. J. Plant Breed, 9(1): 127-145.
- Griffing, B. (1956). concept of general and specific combining ability in relation to diallel crossing system. Aust. J. of Biol. Sci., 9: 463-493.
- Hendawy, F.A.; M.S. Rady; A.M. Abd El-Hamid and R.M. Esmail (1999). Inheritance of fiber traits in some cotton crosses. Egyptian J. Agro., 21: 15-36.

- Khorgade, P.W.; I.V. Satange and L.D. Mmmeshram (2000). Diallel analysis in American cotton (*Gossypium hirsutum* L.). Ind. J. Agric. Res., 34(3): 172-175.
- Laxman, S.M. and Ganesh (2003). Combining ability for yield components and fiber characters in cotton (*Gossypium hirsutum* L.). J. Res. Angra, 31(4): 19-23.
- Rahouma, M.R.A. and S.A. El-Shaarawy (1992). General and specific combining abilities in *G. barbadense*. proc. 5th Conf. Agron. Zagazig 13-15 Sept., 1992, 2: 767-774.
- Zia, U.L.Islam; H.A. Sadaqat and F.A. Khan (2001). Combining ability of some hirsutum cotton types for economic traits. International J. Agric. and Biology, 3(4): 411-412.

التحليل الوراثي للجيل الأول والجيل الثاني في بعض هجن القطن المصري محمد نشأت عبد العزيز نظمي ، أنور عيسى مسعود عيسى ، وليد محمد بسيوني يحيي معهد بحوث القطن – مركز البحوث الزراعية – مصر

خمسة أصناف من القطن تابعة للنوع جوسيبيم باربادنس استخدمت في هذه الدراسة وهي الأصناف جيزة 89 ، جيزة 90 ، جيزة 83، بيما س4 و بيما س6 تم انتخابها وإدخالها في التهجين نصف الدائري وتم الحصول علي الجيل الثاني وزراعته والجيل الأول وذلك لتقدير القدرة العامة والخاصة علي التالف وكذلك درجة التوريث بالمدى الواسع والضيق إلي جانب معاملي التربية الداخلية وذلك لصفات وزن اللوزة و محصول القطن الزهر جم/نبات ، محصول القطن الشعر جم/نبات ، تصافي الحليج ، معاملي البذرة ، طول التيلة بالمليمتر ، المتانة بالجرام تكس وكذلك النعومة بالميكرونير . أظهرت النتائج لقياسات القدرة العامة علي التالف وجود اختلافات معنوية لصفات محصول القطن الزهر و محصول القطن الشعر وطول التيلة في كلا الجيلين وكذلك أيضا لمتانة التيلة وأظهرت قيمة GCA/SCA ان التباين الإضافي كان له السبق والتحكم في توريث صفات وزن اللوزة ، تصافي الحليج ، معاملي البذرة و النعومة في الجيل الأول و لصفات وزن اللوزة ، محصول القطن الشعر ، تصافي الحليج والنعومة في الجيل الثاني .

بالنسبة لتقدير قوة الهجين علي أساس متوسط الآباء كانت معنوية وموجبة للهجين ج 89 x ج 90 لصفات محصول القطن الزهر والشعر والهجين ج 89 x ج 83 لصفات محصول القطن الزهر والشعر وتصافي الحليج والهجين ج 89 x بيما س4 لصفات محصول القطن الزهر ومعاملي البذرة وطول التيلة والهجين ج 89 x بيما س6 لصفات محصول القطن الزهر والشعر وطول التيلة والهجين ج 90 x ج 83 لصفات محصول القطن الزهر والشعر والنعومة ، محصول الزهر والشعر وطول التيلة للهجين ج 90 x بيما س4 ، وطول التيلة والنعومة للهجين ج 83 x بيما س4 والهجين بيما س4 x بيما س6 لصفات محصول القطن الزهر . علي الجانب الآخر بالنسبة لقوة الهجين علي أساس الأب الأفضل أظهرت النتائج وجود اختلافات عالية المعنوية وموجبة للهجين ج 89 x ج 83 لصفة تصافي الحليج ، و صفات معاملي البذرة والطول للهجين ج 89 x بيما س4 ، محصول القطن الزهر والشعر للهجين ج 89 x بيما س6 و ج 90 x ج 83 و ج 90 x بيما س4 علي الجانب الآخر هناك بعض الهجن التي أظهرت قوة هجين منخفضة .

بالنسبة لمعاملي التربية الداخلية كان معنويا وموجبا لصفة محصول القطن الشعر للهجين $(P_1 \times P_2)$ و لصفة وزن اللوزة و محصول القطن الزهر للهجين $(P_1 \times P_3)$ و محصول القطن الزهر ومعاملي البذرة للهجين $(P_1 \times P_4)$ و محصول القطن الزهر والشعر وتصافي الحليج للهجين $(P_1 \times P_5)$ و $(P_2 \times P_4)$ إلي جانب أن معاملي التربية الداخلية كان معنويا وسالب لبعض الهجن تحت الدراسة

قام بتحكيم البحث

أ.د / ممدوح محمد عبد المقصود
كلية الزراعة – جامعة المنصورة
مركز البحوث الزراعية

أ.د / عبد المعطي محمد علي زينة

Table 3: Estimates of specific combining ability effects (S_{ij}) for studied traits.

Traits	Generation	$P_1 \times P_2$	$P_1 \times P_3$	$P_1 \times P_4$	$P_1 \times P_5$	$P_2 \times P_3$	$P_2 \times P_4$	$P_2 \times P_5$	$P_3 \times P_4$	$P_3 \times P_5$	$P_4 \times P_5$	LSD		S_{ij}	$S_{ii} - S_{ij}$
												1%	5%		
B.W.	F ₁	-0.200**	0.105	-0.148**	-0.110*	-0.205**	0.110*	0.048	0.110**	0.019	-0.100	0.143	0.106	0.052	0.090
	F ₂	-0.373**	-0.297**	0.346**	0.203**	0.003	0.313**	0.003	0.313**	-0.021	-0.078	0.156	0.116	0.057	0.098
S.C.Y.	F ₁	5.948	17.286**	7.905	24.157**	12.914**	13.700**	3.019	13.700**	-17.143**	7.576	11.455	8.491	4.146	7.181
	F ₂	6.425	-3.532	-26.765**	-14.494	-19.289*	-11.789	-5.084	-11.789	-24.575**	-6.375	21.694	16.080	7.852	13.600
L.C.Y	F ₁	2.094	9.022**	2.275	11.137**	6.579**	4.865**	-0.440	4.865**	-7.378**	2.208	4.553	3.375	1.648	2.854
	F ₂	-0.484	6.578**	-5.375**	0.059	-2.651	-5.337**	5.497**	-5.337**	-4.808*	2.840	4.851	3.596	1.756	3.041
L.%	F ₁	0.065	1.579**	-0.287	1.179**	0.113	-0.187	-0.854**	-0.187	-0.440	-0.406	0.706	0.523	0.255	0.442
	F ₂	-0.051	0.092	-0.313	-0.222	0.111	-1.894**	-0.570*	-1.894**	0.806**	-0.265	0.658	0.488	0.238	0.412
S.I.	F ₁	-0.517//	0.325**	0.578**	0.116	-0.175	-0.089	0.349**	-0.089	0.059	-0.422**	0.267	0.198	0.097	0.168
	F ₂	-0.162	-0.062	-0.671**	0.524**	0.396**	0.181	-0.190	0.181	-0.157	0.500**	0.282	0.209	0.152	0.177
Length	F ₁	0.571	1.562**	0.943**	0.524	-0.633*	0.948**	-0.571	0.948**	-0.881**	0.367	0.848	0.629	0.307	0.532
	F ₂	0.125	0.211	1.444**	-0.032	0.344	-0.656	0.668	-0.656	0.954**	0.454	0.951	0.705	0.344	0.596
Strength G/tex	F ₁	-0.875	2.383**	-1.151	1.249	-1.622	0.444	-1.522	0.444	-1.598	3.068**	2.347	1.740	0.850	1.471
	F ₂	0.508	0.117	1.441	1.937*	-0.606	-3.149**	0.146	-3.149**	-0.844	1.479	2.079	1.541	0.752	1.303
Mic	F ₁	0.138	0.057	0.543**	-0.005	0.329**	-0.086	-0.200*	-0.086	-0.048	-0.329**	0.259	0.192	0.094	0.163
	F ₂	0.125	0.211	1.444**	-0.032	0.344	-0.656	0.668	-0.656	0.954**	0.454	0.951	0.705	0.344	0.596

*,** significant at 0.05 and 0.01 levels of probability, respectively