

## PROTEOLYSIS AND BIOGENIC AMINES FORMATION IN STERILIZED EDAM CHEESE CURD SLURRY INOCULATED WITH SOME PROBIOTIC STRAINS FORMULATIONS

Rabie, M. A. and A. A. Abdel Galeel

Food Sci. Dept., Fac. Agric., Zagazig Univ., Egypt E-44519

### ABSTRACT

In this investigation, Sterilized Edam cheese curd slurry was inoculated with 1% *Lactococcus lactis subsp. lactis* KF147 and served as control. The effect of added some probiotic strains formulations upon proteolysis pattern and biogenic amines (BAs) during incubation at 30 °C for 21 days. was also evaluated. Sterilized Edam cheese curds slurry were inoculated with 1% of the following formulations: *Propionibacterium shermanii* PS-4 + *Bifidobacterium bifidum* DSM 20082 (1:1), *Propionibacterium shermanii* PS-4 + *Lactobacillus acidophilus* ATCC4356(1:1), or *Propionibacterium shermanii* PS-4 + *Bifidobacterium bifidum* DSM 20082 + *Lactobacillus acidophilus* ATCC4356 (1:1:1). Results showed no significant influence of any formulation of probiotic bacteria on total solids, salt and fat contents of sterilized Edam cheese curd slurry. However, inoculation of sterilized Edam curds slurry with probiotic strains formulations affected the pH and soluble nitrogenous compounds (S.N.C). The formulation consisting of *Propionibacterium shermanii* + *Bifidobacterium bifidum* + *Lactobacillus acidophilus* (1:1:1) showed the highest concentration (7.9 %) of water-soluble nitrogenous compounds at the end of incubation period. While inoculation included *Bifidobacterium bifidum*, a significant reduction in total BAs (447 - 37 mg/kg DW) was observed by 21 d Also, both histamine and tyramine was decreased from 84 - 25, and 359 - 6 mg/kg DW respectively.

**Keywords:** Curd slurry, probiotics, proteolysis, biogenic amines.

### INTRODUCTION

Cheese ripening is a complex process that includes several biochemical reactions leading to cheese flavour and texture development, namely via glycolysis, proteolysis and lipolysis; proteolysis in particular is strongly influenced by the activity of microbial enzymes during ripening (Fox *et al.*, 1993).

Starter cultures account for a major portion of the microflora in young cheese curd; during ripening, starter numbers decrease owing to cell death and subsequent autolysis (Crow *et al.*, 1995). The potential of intracellular peptidases to hydrolyze peptides following autolysis indicates that this may be an important factor in cheese flavour development. Comprehensive experimental evidence suggests that adjunct cultures or non-starter lactic bacteria also play a role in cheese ripening (Peterson *et al.*, 1990; Rodriguez, 1998).

It is of particular interest, the use of cheese as carrier of probiotic strains owing to its mild pH and higher fat content, both of which favour survival when compared to fermented milks (Stratton *et al.*, 1991); this has been consubstantiated for such probiotic species as *Bifidobacterium bifidum*,

*Lactobacillus acidophilus* and *Lactobacillus casei* in curd slurry (Vinderola *et al.*, 2000), as well as *Enterococcus faecium* in yoghurt (Gardiner *et al.*, 1977) – whereas (Vinderola *et al.*, 2002) emphasized the interactions between mesophilic starter and probiotic cultures in cheese.

Biogenic amines (BAs) in cheese have a high impact on public health, since they are potentially toxic to humans at levels  $>100 \text{ mg kg}^{-1}$ . These are low-molecular nitrogenous compounds formed chiefly via microbial decarboxylation of precursor amino acids (Halász *et al.*, 1994). So, they can serve as indicator of food quality Rauscher-Gabernig *et al.*, (2009). Due to the oxidation system encompassing monoamine and diamine oxidases, small amounts of BAs can be metabolized in the organism without any major impact on health – but ingestion of higher levels brings about serious impacts, e.g. release of adrenaline and noradrenaline leading to gastric acid secretion, increased cardiac output, migraine, tachycardia and high blood pressure (Shalaby, 1996; Loret, 2005); furthermore, a fraction of the population has histamine (HIS) intolerance (Maintz *et al.*, 2007).

The ability to decarboxylate amino acids is present in many genera of lactic acid bacteria (LAB) (Halász *et al.*, 1994; Arena *et al.*, 2001) – and tyramine (TYR), HIS, putrescine and cadaverine (CAD) can be found to considerable levels, and other BAs to lesser amounts (Spano *et al.*, 2010). A few strains of lactic acid bacteria, e.g. *Lactococcus lactis* subsp. *lactis*, have exhibited decarboxylase activity (Rabie *et al.*, 2011a); however, lactobacilli have also been associated to release of HIS, TYR and putrescine (PUT), and enterococci to release of TYR (Bover-Cid *et al.*, 1999). The optimum pH of decarboxylases lie usually in the acidic region; therefore, fermented dairy products represent a suitable environment for formation of BAs (Halász *et al.*, 1994; McSweeney *et al.*, 2004) – and *Lactococcus lactis* is presumably responsible for BAs in cheeses at large where it are used as starter culture.

Little data are available on the occurrence of biogenic amines in Edam-type curd slurries. Therefore, the objective of this study was to provide information on the concentration of these components release, by ascertaining the effects of adding probiotic strains to the regular inocula of lactic acid bacteria.

## **MATERIALS AND METHODS**

### **Materials:**

#### **Cow's milk**

Cow's milk, containing 3.5 % fat and 12 % TS, was obtained from Dairy Technology Unit of Food Science Department, Faculty of Agriculture, Zagazig University (Egypt).

#### **Starter and probiotic microorganisms cultures**

*Lactococcus lactis* subsp. *lactis* KF147 and *Propionibacterium shermanii* PS-4 were obtained from Chr. Hansen Laboratories (Copenhagen, Denmark). *Bifidobacterium bifidum* DSM 20082 and *Lactobacillus acidophilus* ATCC4356 were obtained from Cairo Microbiological Center, MIRCEN, Faculty of Agriculture, Ain Shams University (Egypt). Stock cultures were

kept frozen at -70°C in 30 % (v/v) glycerol. Prior to use, each strain was cultivated in MRS broth for 48 h, and underwent two consecutive transfers. When used as adjuncts, the aforementioned strains were inoculated at 0.1% into sterile 2 % fat milk, and incubated for 24 h at the appropriate temperature.

#### **Preparation of Edam cheese curd slurry**

Standardized cow's milk 2% fat was pasteurized at 72°C for 15 sec, cooled to 32°C, and transferred to a sterile cheese vat under a laminar flow hood. Approximately 100 L of milk was used to manufacture Edam cheese curd, following the procedure described by Scot (1979).

Cheese curd slurry was prepared as described by Farkye et al. (1995); it was salted at a rate of 2 %, and transferred aseptically into a sterile, wide mouth bottle, and sterilized at 121°C for 15 min. All aseptic cheese slurries were aseptically inoculated with 1 % of Edam cheese starter. One part of cheese slurry without added probiotic stains served as control. The other portions of cheese slurry were aseptically inoculated with 1% of each combination of *Propionibacterium shermanii* PS-4 + *Bifidobacterium* DSM 20082, or *Propionibacterium shermanii* PS-4 + *Lactobacillus acidophilus* and or *Propionibacterium shermanii* PS-4 + *Lactobacillus acidophilus* ATCC4356 + *Bifidobacterium* DSM 20082 (1:1:1). Each slurry preparation was replicated three times, using freshly made Edam cheese curd slurry. Cheese slurry bottles were incubated at 30°C for 7, 14 and 21 days.

#### **METHODS:**

##### **Assessment of proteolysis**

The pH 4.6-water soluble nitrogen (WSN), 12% trichloroacetic acid soluble nitrogen (TCA-SN), Non protein nitrogen (NPN) and Phosphotungstic acid soluble nitrogen PTA-SN amino acid nitrogen (AN) were determined as described by (Gripon *et al.*, 1975).

##### **Evaluation of flavour intensity**

Flavour intensity of Edam cheese curd slurry with or without probiotic bacteria was evaluated as described by (King *et al.*, 1979).

##### **Extraction of biogenic amines**

Biogenic amines in curd slurry samples were extracted as originally described by (Simon-Sarkadi and Holzaphel., 1994), and later improved by (Rabie *et al.*, 2011b). To extract BAs, 10 mL of 10 % trichloroacetic acid (TCA) was added to 3 g-samples of cheese, and the mixture was shaken for 1 h using a Laboshake (Gerhardt Ls 500i, Germany). The extract was filtered through Whatman No.1 filter paper. To remove the fat content, the samples were kept at -20 °C for 1 day, and then centrifuged (MLW, T 24, Germany) at 7000 xg for 15 min. Supernatants were collected and filtered through a 0.25-µm membrane filter (Nalgene, USA).


##### **Quantification of biogenic amines**

The analysis of BAs was performed with an AAA 400 amino acid analyser (Ingos, Czech Republic), equipped with an Ostion LG ANB ion-exchange column. Colorimetric detection was accomplished at 570 and 440 nm, after post-column derivatization with ninhydrin (Csomos *et al.*, 2002). The operating parameters used were summarized at Table 1.

**Table 1: Analytical conditions utilized for quantitation of biogenic amines.**

Items	Feature	Value
Column	length	7 cm
	diameter	0.37 cm
	temperature	60°C
Eluent	flow rate	0.30 mL min <sup>-1</sup>
	run time	101 min
Reagent	flow rate	0.20 mL min <sup>-1</sup>
	temperature	121 °C
Injection	sample volume	100 µL
Detection	wavelength	570 nm
		A: 6.00
Citric buffer sequence	pH	B: 5.70
		C: 5.45

Calibration curves were prepared in advance with biogenic amine chromatographic standards (see Fig. 1), based on means of triplicate data. Identification and quantification of each biogenic amine were accomplished by comparing retention time and peak area, respectively, between actual samples and standards.



**Fig 1: Amino acid analyzer chromatogram of biogenic amines standard**  
 \*1. Histamine 2. Tyramine 3. Putrescine 4. Cadaverine 5. Spermidine 6. Agmatine  
 7. Spermine

**Statistical analyses**

Statistical analysis of the data produced was via one-way ANOVA and Tukey’s test; statistical significance was declared at 5%.

**RESULTS AND DISCUSSION**

The average chemical composition of Edam cheese curd slurries containing probiotic bacterial strains, during ripening at 30°C, is presented in Table 2. Addition of probiotic bacterial strains apparently did not influence chemical composition. Similar results were reported by (Karakus *et al.*, 1995; Mara *et al.*, 1999; Ong *et al.* 2007).

Changes in pH, WSN, TCA-SN and PTA-SN will hereafter be taken as indices of ripening of Edam-type curd slurry. The pH of cheese curd slurries ranged from 5.60 to 5.15, and pH in slurries containing probiotic strains was slightly lower than that of the control. This could be explained by development of acidity at high incubation temperature (30°C) when added with probiotic bacterial strains.

**Table 2: Effect of combination of probiotic strains upon chemical composition of Edam cheese curd slurry during incubation.\***

Item %	Incubation period (days)											
	7				14				21			
	C	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)	C	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)	C	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)
Total Solids	40.10	40.20	40.33	40.50	40.20	40.35	40.25	40.30	40.05	40.20	40.33	40.30
Salt	1.78	1.76	1.72	1.84	1.58	1.86	1.92	1.68	1.64	1.76	1.72	1.94
Fat	12.59	12.66	12.82	12.91	12.70	12.79	12.80	12.85	12.64	12.72	12.80	12.84

\*C = Control Edam cheese curd containing *L. lactis subsp. lactis*  
 P+B = Propionibacterium shermanii + Bifidobacterium bifidum (1:1)  
 P+L = Propionibacterium shermanii + Lactobacillus acidophilus (1:1)  
 P+B+L = Propionibacterium shermanii + Bifidobacterium bifidum + Lactobacillus acidophilus (1:1:1)

The curd slurry containing *B. bifidum* DSM 20082 + *L. acidophilus* ATCC4356 + *P. shermanii* PS-4 strains (1:1:1) exhibited the highest ( $P<0.05$ ) concentration of WSN (Table 3). Kasimoglu *et al.*, (2004) reported that *L. acidophilus* produced an increase in this fraction during ripening of probiotic white cheese.

The levels of WSN, TCA-SN and PTA-SN increased in all samples during incubation as a consequence of proteolysis (Table 3) – but significant differences ( $P<0.05$ ) between cheeses with and without added probiotic bacteria were noticeable.

**Table 3: Effect of combinations of probiotic strains upon soluble nitrogen fractions of Edam cheese curd slurry during incubation.\***

Nitrogen fractions (% of TN)	Incubation period (days)											
	7				14				21			
	C*	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)	C*	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)	C*	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)
WSN/TN**	5.40a	7.98b	9.8c	10.6c	7.75a	10.87b	12.5c	13.88d	10.89a	12.88b	14.76c	16.77d
NPN/TN**	3.15a	4.35b	4.99c	5.39d	4.10a	4.90b	5.88c	6.55d	5.87a	7.44b	7.34b	7.90b
AN/TN**	2.25a	3.66b	4.44c	5.27d	3.88a	5.66b	6.88c	7.86d	4.66a	5.44b	6.25c	7.88d
pH	5.60a	5.45a	5.25b	5.15b	5.60a	5.35b	5.30b	5.00c	4.70a	4.75a	4.90b	4.95b

\*C = Control Edam cheese curd containing *L. lactis subsp. lactis*  
 P+B = Propionibacterium shermanii + Bifidobacterium bifidum (1:1)  
 P+L = Propionibacterium shermanii + Lactobacillus acidophilus (1:1)  
 P+B+L = Propionibacterium shermanii + Bifidobacterium bifidum + Lactobacillus acidophilus (1:1:1)

The value of WSN in the control increased 2.3 fold by the end of incubation, and the evolution in WSN, TCA-SN and PTA-SN was similar to

that reported by (Macedo *et al.*, 1997). The obtained results indicate that probiotic bacteria present a significant effect upon secondary proteolysis. In addition, Edam-type slurry inoculated with *B. bifidum* DSM 20082 + *L. acidophilus* ATCC4356 + *P. shermanii* PS-4 strains (1:1:1) showed the highest ( $P<0.05$ ) concentration of WSN (16.8 %) and TCA-SN (7.8 %). These results resemble those of Madkor *et al.*, (2000), who reported that adjunct cultures of similar strains of lactobacilli can increase the level of PTA-SN. Whereas, Drake *et al.* (1996) found that *Lactobacillus helveticus*, used as adjunct starter in reduced fat Cheddar, also produces significantly greater rates of proteolysis than the control.

The results on flavour intensity in assessment between control and probiotic Edam cheese curd slurries are shown in Table 4.

**Table 4: Effect of combinations of probiotic strains upon development of flavour intensity of Edam cheese curd slurry during incubation.\***

Incubation period	Probiotic strains combinations											
	C	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)	C	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)	C	P+B (1:1)	P+L (1:1)	P+B+L (1:1:1)
7	4.40	4.65	4.85	4.90	4.95	5.35	5.5	5.88	5.30	5.60	5.80	5.88
14	4.75	4.85	4.99	5.10	5.1	5.9	5.88	5.55	5.60	5.44	5.90	5.90
21	4.75	5.50	5.74	5.87	5.20	5.30	5.45	5.65	5.65	5.74	5.85	5.92

\*C = Control Edam cheese curd containing *L. lactis* subsp. *lactis*


P+B = *Propionibacterium shermanii* + *Bifidobacterium bifidum* (1:1)

P+L = *Propionibacterium shermanii* + *Lactobacillus acidophilus* (1:1)

P+B+L = *Propionibacterium shermanii* + *Bifidobacterium bifidum* + *Lactobacillus acidophilus* (1:1:1)

Addition of *P. shermanii* + *L. acidophilus* (1:1) or *P. shermanii* + *B. bifidum* (1:1) or *P. shermanii* + *L. acidophilus* + *B. bifidum* (1:1:1) significantly improved the flavour of Edam cheese curd slurries; this could be attributed to the higher concentration of soluble nitrogenous compounds.

Five biogenic amines, viz. histamine, tyramine, putrescine, spermine and spermidine were found in Edam curds slurries; both control and experimental ones (see Fig 2, 3, and 4). The control Edam curd slurry (Fig 2.) showed the highest concentration of total BAs (478.84 mg/kg DW by 21 days), which almost doubled between 7 and 21 days.



**Fig. 2 Changes in the contents of total biogenic amines in Edam cheese curd slurry during incubation.**

\*C = Control Edam cheese curd containing *L. lactis* subsp. *lactis*


P+B = *Propionibacterium shermanii* + *Bifidobacterium bifidum* (1:1)

P+L = *Propionibacterium shermanii* + *Lactobacillus acidophilus* (1:1)

P+B+L = *Propionibacterium shermanii* + *Bifidobacterium bifidum* + *Lactobacillus acidophilus* (1:1:1)

The lowest content (5.2-35.1 mg/kg DW) of total BAs was observed in Edam-type curd slurry inoculated with *P. shermanii* + *B. bifidum* + *L. acidophilus* (1:1:1), followed closely (35.2-63.1 mg/kg DW) by *P. shermanii* + *B. bifidum* (1:1), and finally by *P. shermanii* + *L. acidophilus* (1:1) (46.8-74.84 mg/kg DW) as illustrated in Fig. 2. Similar results in terms of proteolytic breakdown were reported by Komprda *et al.*, (2007a) and Komprda *et al.*, (2008b). A rapid pH decrease may prevent BA formation in fermented products; and starter cultures able to outgrow nonstarter bacteria during late ripening and storage may also avoid excessive BA buildup (Suzzi *et al.*, 2003) – for instance, Roig-Sagués *et al.*, (1999) claimed that *Lactobacillus sake* is more suitable than *Lactobacillus curvatus* as starter culture if low BA levels is used as processing goal.

The changes in concentration of HIS are depicted in Fig. 3; this BA was found at high content 148.2 mg/kg Dw in non-inoculated Edam cheese – which is 3-fold the recommended maximum level (EFSA, 2011) by 21 days.



**Fig. 3 Changes in the contents of histamine, tyramine and putrescine in Edam cheese curd slurry during incubation**

\*C = Control Edam cheese curd containing *L. lactis* subsp. *lactis*

P+B = *Propionibacterium shermanii* + *Bifidobacterium bifidum* (1:1)

P+L = *Propionibacterium shermanii* + *Lactobacillus acidophilus* (1:1)

P+B+L = *Propionibacterium shermanii* + *Bifidobacterium bifidum* + *Lactobacillus acidophilus* (1:1:1)


A reduction of HIS was observed with *P. shermanii* + *B. bifidum* + *L. acidophilus* (1:1:1); its concentration was lower than the control 238 and the other probiotic bacterial combinations tested.

The content of TYR remained high in non-inoculated cheese samples, and ranged from 190.4 to 359.2 mg/kg DW during the ripening period (see Fig. 3). On the other hand, addition of some strain combinations dramatically reduced formation of TYR during ripening: *P. shermanii* + *B. bifidum* + *L. acidophilus* (1:1:1) was more effective in this respect. Priyadarshani et al. (2011) showed that BA reduction was not apparent for *L. acidophilus*, *L. lactis* subsp. *lactis*, *L. lactis* subsp. *lactis* and *Lactobacillus plantarum*; they claimed that BA formation is strain dependent but not related only to species. Therefore, careful screening for amino acid decarboxylase activity is recommended before selecting LAB or probiotic strains as starter in dairy industry (Komprda et al 2007a; Komprda et al 2008b; Priyadarshani et al., 2011). Similar results were obtained by (Bunková et al., 2010) for selected biogenic amines (HIS, TYR, and PUT) in 4 layers of Dutch-type cheese (Edam-cheese), depending on the ripening/ storage regime followed along cheese ripening.

The combination consisting of *P. shermanii* + *B. bifidum* + *L. acidophilus* (1:1:1) was more pronounced in the reduction of PUT; it increased from 12.8 mg/kg DW by 7 days to a 2.6-fold value by 21 days of ripening (see Fig. 3). This result in a dairy matrix is consistent with the reduction in putrescine in sauerkraut (plant matrix) inoculated with *Lactobacillus plantarum*, *L. casei* and *Lactobacillus curvatus* (Rabie et al., 2011a) and in sukuc (meat matrix) inoculated with *Lactobacillus sakei* and *Staphylococcus carnosus* (Genççelep et al., 2007) – so acidity may play a



role. Finally, the maximum contents of spermidine and spermine in non-inoculated Edam cheese were 3.1 and 3.6 mg/kg DW, as shown in Fig. 4. There is also a decrease of 1.6 and 2.1-fold of the concentration of spermidine and spermine, especially in the case of Edam cheese inoculated with *P. shermanii* + *B. bifidum* + *L. acidophilus* (1:1:1). These two amines accounted for the lowest concentrations.



**Fig. 4 Changes in the contents of spermidine and spermine in Edam cheese curd slurry during incubation.**

\*C = Control Edam cheese curd containing *L. lactis* subsp. *lactis*

P+B = *Propionibacterium shermanii* + *Bifidobacterium bifidum* (1:1)

P+L = *Propionibacterium shermanii* + *Lactobacillus acidophilus* (1:1)

P+B+L = *Propionibacterium shermanii* + *Bifidobacterium bifidum* + *Lactobacillus acidophilus* (1:1:1)

The unique performance of the co-cultures added containing the two probiotic strains (i.e. *B. bifidum* and *L. acidophilus*) arises likely from their absence of the enzymatic machinery required to decarboxylate free amino acids, coupled with a putative capacity to actually take up existing BAs; their ecological dominance in the curd slurry will eventually hamper net formation of biogenic amines, and even actively contribute to reduce the current level of BAs.

## REFERENCES

- Arena, M.E. and Manca de Nadra, M.C. (2001). Biogenic amine production by *Lactobacillus*. *Journal of Applied Microbiology*, 90, 158–162.
- Bover-Cid, S. and Holzapfel, W. H. (1999). Improved screening procedure for biogenic amine production by lactic acid bacteria. *International Journal of Food Microbiology*, 53, 33–41.
- Bunková, L.; Bunka, F.; Mantlová, G.; Cablová, A.; Sedláček, I. Svec, P.; Pachlová, V. and Krácmár, S. (2010). The effect of ripening and storage conditions on the distribution of tyramine, putrescine and cadaverine in Edam-cheese. *Food Microbiology*, 27, 880-888.
- Crow, V. L., T. Coolbear, P. K. Gopal, F. G. Martley, L. L. McKay, and H. Riepe. (1995). The role of autolysis of lactic acid bacteria in the ripening of cheese. *International Dairy Journal*, 5, 855–875.

- Csomos, E., and Simon-Sarkadi, L. (2002). Characterisation of tokaj wines based on free amino acid and biogenic amine using ion-exchange chromatography. *Chromatographia*, 56, 185–188.
- Drake, M. A., T. D. Boylston, K. D. Spence, and B. G. Swanson. (1997). Improvement of sensory quality of reduced fat Cheddar cheese by *Lactobacillus adjunct*. *Food Research International*, 30, 35–40.
- EFSA – European Food Safety Authority; Panel on Biological Hazards (BIOHAZ) (2011). Scientific opinion on risk base control of biogenic amine formation in fermented foods. *EFSA Journal*, 9, 2393–2486.
- Farkye, N. Y.; Madkor, S. A. and Atkins, H. G. (1995). Proteolytic abilities of some lactic acid bacteria in a model cheese system. *International Dairy Journal*, 5:715 –725.
- Fox P F, Law J, McSweeney P L H and Wallace J (1993) Biochemistry of cheese ripening. In *Cheese: Chemistry, Physics and Microbiology*, Vol 1: General Aspects, 2nd edn, pp 389–438. Fox P F, ed. London: Chapman & Hall.
- Gardiner, G., Stanton, C., Lynch, P.B., Collins, J.K., Fitzgerald, G., and Ross, R.P. (1999). Evaluation of Cheddar cheese as a food carrier for delivery of a probiotic strain to the gastrointestinal tract. *Journal of Dairy Science*, 82, 1379-1387.
- Genççelep, H., Kaban, G. and Kaya, M. (2007). Effect of starter cultures and nitrite levels on formation of biogenic amines in sucuk. *Meat Science*, 77, 424–430.
- Gripon, J. C.; Desmazeaud, M. J.; Bars, D. L. and Bergre, L. (1975). Study of the role of microorganisms and enzymes during ripening of cheeses. II. Influence of commercial rennin. *Le Lait*, 55,502-509.
- Halász, A., Baráth, Á., Simon-Sarkadi, L., Holzapfel, W. (1994). Biogenic amines and their production by microorganisms in food. *Trends Food Science and Technology* 51, 42-49.
- Karakus M., Alperden I. (1995). Effect of starter composed of various species of lactic acid bacteria on quality and ripening of Turkish White pickled cheese, *Lebensmittel-Wissenschaft & Technologie*. 28:404–409.
- Kasimoglu, A., Göncüglü, M., Akgün S. (2004). Probiotic white cheese with *Lactobacillus acidophilus*. *International Dairy Journal*, 14, 1067-1073.
- King, R. D. and Clegg, G. H. (1979). The metabolism of fatty acids, methyl ketones and secondary alcohols by *Penicillium roquefortii* in blue cheese slurries. *Journal of Science Food Agriculture*, 30, 197-202.
- Komprda, T., Smelá, D., Novická, K., Kalhotka, L., \_Sustová, K., Pechová, P., (2007a). Content and distribution of biogenic amines in Dutch-type hard cheese. *Food Chemistry*, 102, 129-137.
- Komprda, T., Burdychová, R., Dohnal, V., Cwíková, O., Sládková, P., Dvorácková, H., (2008b). Tyramine production in Dutch-type semi-hard cheese from different producers. *Food Microbiol.* 25, 219-227.
- Loret, S., Deloyer, P. and Dandrifosse, G. (2005). Levels of biogenic amines as a measure of the duality of the beer fermentation process: data from Belgian samples. *Food Chemistry*, 89, 519–525.

- Macedo, A. C., and Malcata, F. X. (1997). Secondary proteolysis in Serra cheese during ripening and throughout the cheesemaking season. *Zeitschrift für Lebensmittel Untersuchung und Forschung A*, 204, 173–179.
- Madkor, S.A.; Tong, P.S. and El Soda, M (2000). Ripening of Cheddar cheese with attenuated adjunct cultures of lactobacilli. *Journal of Dairy Science*, 83, 1684-1691.
- Maintz, L. and Novak, N. (2007). Histamine and histamine intolerance. *American Journal of Clinical Nutrition*, 85, 1185–1196.
- Mara, O., and Kelly A.L. (1999). Contribution of milk enzymes, starter and rennet to proteolysis during storage of Quarg, *International Dairy Journal*, 8, 973–979.
- McSweeney, P. (2004). Biochemistry of cheese ripening. *International Journal of Dairy Technology*, 57, 127–144.
- Ong, L.; Henriksson, A. and Shah, N. P. (2007). Proteolytic pattern and organic acid profiles of probiotic Cheddar cheese as influenced by probiotic strains of *Lactobacillus acidophilus*, *Lb. paracasei*, *Lb. casei* or *Bifidobacterium* sp. *International Dairy Journal*, 17, (1), 67-78.
- Peterson, B. D. and R. T. Marshall. (1990). Non-starter lactobacilli in Cheddar cheese: A review. *Journal of Dairy Science*, 73:1395–1410.
- Priyadarshani, W. M. D. and Rakshit, S.K. (2011): Screening selected strains of probiotic lactic acid bacteria for their ability to produce biogenic amines (histamine and tyramine). *International Journal of Food Science & Technology*, 46, 10, 2062–2069.
- Rabie, M.A., Siliha, H.I., El-Saidy, S.M, El-Badawy, A.A., and Malcata, F.X. (2011a). Reduced biogenic amine contents in sauerkraut via addition of selected lactic acid bacteria. *Food Chemistry*, 129, 1778–1782.
- Rabie M.A., Siliha, H.I., El-Saidy, S.M, El-Badawy, A.A., Malcata F. X. (2011b). Effect of -irradiation upon biogenic amine formation in blue cheese during storage *International Dairy Journal*, 21, 373-376.
- Rauscher-Gabernig, E., Grossgut, R., Bauer, F. and Paulsen, P. (2009). Assessment of alimentary histamine exposure of consumers in Austria and development of tolerable levels in typical foods. *Food Control*, 20, 423–429.
- Rodriguez, J. (1998). Recent advances in the development of low-fat cheeses. *Trends in Food Science & Technology*, 9, 249–254.
- Roig-Sagués, A.X., and Eerole, S. (1999). Biogenic amines in meat inoculated with *Lactobacillus* sake starter strains and an amine-positive lactic acid bacterium. *Zeitschrift für Lebensmittel Untersuchung und Forschung A*, 205, 227–231.
- Scott, R. (1981): Cheese making practice. Applied Science Publishers Ltd London.
- Shalaby, A.R. (1996). Significance of biogenic amines to food safety and human health. *Food Research. International*, 29, 675-690.
- Simon-Sarkadi, L., and W. Holzapfel. (1994). Determination of biogenic amines in leafy vegetables by amino acid analyser. *Zeitschrift für Lebensmittel Untersuchung und-Forschung A*, 198, 220–233.

- Spano, G., Russo, P., Lonvaud-Funel, A. (2010): Biogenic amines in fermented foods. *European Journal of Clinical Nutrition*, 64, S95–S100.
- Stratton, J.E., Hutkins, R.W, Taylor, S.L. (1991). Biogenic amines in cheese and other fermented foods: a review. *Journal of Food Protection*, 54, 460–470.
- Suzzi, G., and Gardini, F. (2003). Biogenic amines in dry fermented sausages: a review. *International Journal of Food Microbiology*, 88, 41–54.
- Vinderola, C.G, Prosello, W., Ghiberto, D., and Reinheimer, J.A. (2000). Viability of probiotic (*Bifidobacterium*, *Lactobacillus acidophilus* and *Lactobacillus casei*) and nonprobiotic microflora in Argentinian Fresco cheese. *Journal of Dairy Science*, 83, 1905–1911.
- Vinderola, C.G., Mocchiutti, P., and Reinheimer, J.A. (2002). Interactions among lactic acid starter and probiotic bacteria used for fermented dairy products. *Journal of Dairy Science*, 85, 721–729.

**التحلل البروتيني وتكوين الأمينات الحيوية في معلق جبن الايدم المعقمة و الملحقة ببعض التوليفات من سلالات البكتيرية الحيوية**  
**محمد عبدالحميد ربيع و علي عبد الرحمن عبد الجليل**  
**قسم علوم الأغذية-كلية الزراعة-جامعة الزقازيق**

في هذا البحث تم تلقیح معلق الخثرة المعقمة لجبن الايدام (المقارنة) بنسبة ١٪ من بكتريا *Lactococcus lactis subsp. lactis* KF147 ولدراسة تأثير اضافة بعض التوليفات من سلالات البكتيري الحيوية على التحلل البروتيني وتكوين الأمينات الحيوية اثناء التحضين على درجة ٣٠م لمدة ٢١ يوم فقد تم تلقیح معلق الخثرات المعقمة لجبن الايدام بنسبة ١٪ من التوليفات الأتية: *Propionibacterium shermanii* PS-4 + *Bifidobacterium bifidum* DSM 20082 (1:1), *Propionibacterium shermanii* PS-4 + *Lactobacillus acidophilus* (1:1) ATCC4356, or *Propionibacterium shermanii* PS-4 + *Bifidobacterium bifidum* DSM 20082 + *Lactobacillus acidophilus* ATCC4356 (1:1:1). وأظهرت النتائج عدم وجود تأثير معنوي من أي توليفة من البكتريا الحيوية على المواد الصلبة الكلية والملح والدهن لمعلق الخثرات المعقمة لجبن الايدام. ومع ذلك، فقد وجد ان تلقیح الخثرات المعقمة لجبن الايدام قد أظهر تأثيرا على رقم الحموضة والمركبات النيتروجينية الذائبة كما أظهرت التوليفة التي تتكون من *Propionibacterium shermanii* + *Bifidobacterium bifidum* + *Lactobacillus acidophilus* (1:1:1) أعلى تركيز (٧.٩٪) من النيتروجين الذائب في الماء في نهاية فترة التحضين. وعندما اشتمل التلقیح على *Bifidobacterium bifidum* لوحظ انخفاض كبير في تركيز الأمينات الحيوية الكلية (٤٤٧-٣٧ ملليجرام / كجم بالنسبة للوزن الجاف) كما لوحظ ايضا انخفاض في تركيز الهستامين (٨٤-٢٥) والتيرامين (٣٥٩ - ٦ ملليجرام / كجم بالنسبة للوزن الجاف) على التوالي.